VOL. 251 JULY 1953 No. 7

# American Journal of Science

Established in 1818 by Benjamin Silliman

EDITORS
CHESTER R. LONGWELL - JOHN RODGERS

ASSOCIATE EDITORS

REGINALD A. DALY, CAMBRIDGE, MASS.
A. F. BUDDINGTON, PRINCETON, N. J.
CARL O. DUNBAR
HENRY MARGENAU
HORACE WINCHELL
RICHARD F. FLINT
HAROLD G. CASSIDY
G. EVELYN HUTCHINSON
NEW HAVEN, CONN.
LEASON H. ADAMS
WILMOT H. BRADLEY
WASHINGTON, D. C.
WILLIAM H. TWENHOFEL, ORLANDO, FLORIDA
HOWEL WILLIAMS, BERKELEY, CALIF.
FREDERICK J. ALCOCK, OTTAWA, CANADA
CECIL E. TILLEY, CAMBRIDGE, ENGLAND
GEORGE GAYLORD SIMPSON, NEW YORK, N. Y.
ADOLPH KNOPF, STANFORD, CALIF.

STERLING TOWER
NEW HAVEN, CONNECTICUT

Published monthly. Eight dollars a year. Seventy-five cents a copy.
Foreign, eight dollars sixty cents. Canada, eight dollars thirty-five cents.

# AMERICAN JOURNAL OF SCIENCE

*Established in 1818 by Benjamin Silliman*

THE FIRST SCIENTIFIC JOURNAL IN THE UNITED STATES

DEVOTED TO THE GEOLOGICAL SCIENCES AND TO RELATED FIELDS

*Editor*: CHESTER RAY LONGWELL
*Assistant Editor*: JOHN RODGERS
*Executive Secretary*: ELIZABETH G. WEINMAN

Entered as second-class matter at the Post Office at New Haven, Conn., under the Act of March 3, 1879. Published monthly at Sterling Tower, Yale University, New Haven, Conn.

*Subscription rate* $8 per year (75 cents a number). Student rate $5 per year (application blanks available on request). Postage prepaid to the United States and Central and South America; 35 cents per year to Canada; 60 cents per year elsewhere.

The JOURNAL completed its *first* series of 50 volumes in 1845, its *second* series of 50 volumes in 1870, its *third* series of 50 volumes in 1895, its *fourth* series of 50 volumes in 1920, its *fifth* series of 36 volumes in 1938. Since 1938 the numbering by series has been discontinued, and volumes are annual and bear the whole numbers as of 1818, that for 1939 being volume 237.

Back numbers and volumes, also cumulative indices of all series, may with few exceptions be obtained from the office of the AMERICAN JOURNAL OF SCIENCE, Sterling Tower, New Haven, Conn. Positive microfilm copies of each complete volume may be purchased by subscribers to the paper edition from *University Microfilms*, 313 N. First St., Ann Arbor, Michigan. Regular subscribers may purchase each past year in microcard form, from *The Microcard Foundation*, Wesleyan University, Middletown, Connecticut.


## INSTRUCTIONS TO CONTRIBUTORS

*Manuscripts* of articles submitted should follow the recommendations in the pamphlet "Suggestions to Authors" (Lane, 1935) and should be preceded by brief abstracts. They should be checked with special care before being submitted, to avoid changes in proof other than printer's errors.

*References to literature* should be listed alphabetically by authors at the end of the article. All references should be complete and should comprise in order: Author's name, year of publication, full title: name of periodical or series, volume number, inclusive pages, place of publication (if needed to identify book or periodical). Except that the year should directly follow the author's name, references should follow the form given in "Suggestions to Authors" (Lane, 1935, p. 16-29). As in this example, citations within the article should be to author and year, and to specific pages if appropriate.

*Plates* (photographs), *figures* (line drawings), and *footnotes* should each be numbered consecutively through each article, using arabic numerals. If two photographs form one plate, they are figures 1 and 2 of that plate.

*Illustrations* should be neat and legible and should include explanations of symbols used. Copy that cannot be reproduced cannot be accepted. Plates and figures should be capable of reduction to not more than 4 by 7 inches, all lettering being at least 1/16th inch high after reduction. When necessary, one large map or table can be accepted, if it will not exceed 7 inches in width after reduction. Line drawings should be in black India ink on white drawing board, tracing cloth, or coordinate paper printed in blue. Photographs should be positive glossy prints. Photostatic and typewritten material cannot be accepted as copy for illustrations.

*Reprints.* Thirty separate copies of each article will be furnished to the author author free of cost and without previous notice from him; these will be without a cover. Additional copies will be furnished at cost, which will of course be greater if the article is accompanied by plates involving unusual expense. Copies will be furnished with a printed cover giving the title, author, volume, page, and year, when specially ordered.

### REFERENCE


Lane, B. H., 1935, Suggestions to authors of papers submitted for publication by the United States Geological Survey: U. S. Geol. Survey, Washington, D. C. (For sale by Superintendent of Documents, Washington, D. C., 35c.)

[American Journal of Science, Vol. 251, July 1953, Pp. 481-504]

# American Journal of Science

*JULY 1953*

## PETROLOGICAL CALCULATIONS IN METASOMATIC PROCESSES

ARIE POLDERVAART

ABSTRACT. Petrological calculations in metasomatic processes are critically reviewed. The six analyses of the Stavanger metamorphic series of Norway are used as examples in different methods of calculation. The application of the so-called law of equal volume to all cases of metasomatic changes in composition is questioned. A new method is suggested in which the number of (Si,Al)$O_4$ tetrahedra is assumed to remain constant. In terms of mineralogy and petrology this is believed to have more meaning than the assumption of constant volume.

### INTRODUCTION

METASOMATISM has been defined by Lindgren (1933, p. 91) as "the process of practically simultaneous . . . solution and deposition by which a new mineral of partly or wholly differing . . . composition may grow in the body of an old mineral or mineral aggregate." It follows that metasomatic processes take place in the presence of a fluid phase—water—which acts as a solvent. The amount of pore fluid present in metasomatism is believed to vary, from the small fraction represented by "the intergranular film" to ten per cent or more of the total volume. Lindgren also distinguishes between (1) metasomatic changes proceeding in free crystals or loose aggregates where the force of crystallization can easily overcome the retaining pressure, and (2) metasomatic changes proceeding in rigid rocks where the new mineral makes room for itself by solution of the host mineral. In the first case there is a relatively large amount of pore fluid and metasomatism is accompanied by volume changes, while in the second case the amount of pore fluid is very small and the volume remains constant. The so-called law of equal volume has become the basis for all petrological calculations of metasomatic changes in bulk composition, even when there is

good reason to believe that changes in volume did occur during the process. Such calculations are especially important in determining compositional changes in metamorphism and granitization, i.e., for rocks which consist predominantly of silicates.

In this paper the various methods of calculation are reviewed particularly as applied to silicate rocks. Silicates possess a special type of structure in which (Si, Al)$O_4$ tetrahedra act as fundamental building blocks. Different structures and hence different minerals result from a different arrangement of these blocks and the occupation of the interstitial spaces by different cations. Thus it is suggested that a more realistic approach to the problem is obtained by assuming that during metasomatism of silicate rocks the number of (Si, Al)$O_4$ tetrahedra remains approximately constant.

The classic metamorphic series of the Stavanger region of Norway (Goldschmidt, 1920) is used throughout as example, also because this series has been quoted repeatedly in the recent literature (Reynolds, 1946; Chao, 1951; Joplin, 1952). Table 1 gives the original data. These have been recalculated to 100 per cent after subtraction of certain constituents listed in table 2,[1] while the specific gravity values have also been corrected. $H_2O^-$ has been subtracted, but $H_2O^+$ is retained in the recalculated analyses, the assumption being that $H_2O^+$ represents water chemically bound in the rockforming minerals. There is good reason to believe that this is not true (Faber, 1941), but probably the error in thus regarding $H_2O^+$ is less than that introduced in recalculating the analyses on an anhydrous basis. This emphasizes the great need for a chemical method whereby $H_2O$ which enters into the composition of rockforming minerals can be accurately determined and distinguished from absorbed water. The results are given in table 3, and these data are used throughout in the following calculations.

## CALCULATIONS BASED ON CONSTANT VOLUME

In the most generally used method of calculating changes in composition during metasomatism the analyses are recal-

[1] Subtraction of these constituents is of course not essential, but this has been done to give a more uniform and simplified statement of the six analyses.

TABLE 1

The Stavanger Metamorphic Series; Original Data

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $SiO_2$ | 58.32 | 57.29 | 60.62 | 62.40 | 64.35 | 64.70 |
| $TiO_2$ | .98 | .97 | .80 | .81 | .70 | .59 |
| $Al_2O_3$ | 20.00 | 20.60 | 18.50 | 16.27 | 16.54 | 15.45 |
| $Fe_2O_3$ | 2.01 | 2.40 | 1.32 | 1.44 | 2.24 | 1.36 |
| FeO | 4.98 | 5.48 | 5.65 | 5.78 | 3.92 | 4.18 |
| MnO | .22 | .41 | .07 | .35 | .14 | .05 |
| MgO | 1.85 | 1.76 | 2.42 | 2.19 | 2.04 | 1.48 |
| CaO | .66 | .37 | 1.66 | 2.07 | 2.04 | 2.92 |
| BaO | .10 | .10 | .... | .07 | .11 | .10 |
| $Na_2O$ | 1.26 | 1.40 | 1.81 | 2.28 | 2.00 | 3.09 |
| $K_2O$ | 4.49 | 4.30 | 4.02 | 3.16 | 3.82 | 3.46 |
| $H_2O^+$ | 4.05 | 4.00 | 2.92 | 2.25 | 1.73 | 1.82 |
| $H_2O^-$ | .05 | .05 | .17 | .16 | .12 | .15 |
| $P_2O_5$ | .15 | .14 | .08 | .16 | .17 | .19 |
| $CO_2$ | .43 | .05 | .... | .52 | .13 | .85 |
| C | .36 | .81 | .... | .10 | .... | .... |
| S | .06 | .11 | .05 | .04 | .... | .05 |
| $Cr_2O_3$ | .02 | .... | .... | .... | .... | .... |
| $V_2O_3$ | .03 | .... | .... | .... | .... | .... |
| | 100.02 | 100.24 | 100.09 | 100.05 | 100.05 | 100.44 |
| Sp. gr. | 2.798 | 2.862 | 2.838 | 2.814 | 2.812 | 2.777 |

1. Quartz-muscovite-chlorite phyllite.
2. Quartz-muscovite-chlorite-garnet phyllite.
3. Quartz-muscovite-biotite-garnet phyllite.
4. Quartz-muscovite-biotite-garnet schist.
5. Quartz-muscovite-biotite-garnet mica-schist.
6. Albite porphyroblast schist.

TABLE 2

Material Subtracted in Recalculating Analyses

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $FeS_2$ | .13 | .21 | .11 | .08 | .. | .11 |
| $CaCO_3$ | .93 | .11 | .. | 1.18 | .30 | 1.93 |
| $H_2O^-$ | .05 | .05 | .17 | .16 | .12 | .15 |
| C | .36 | .81 | .. | .10 | .. | .. |
| $Cr_2O_3$ | .02 | .. | .. | .. | .. | .. |
| $V_2O_3$ | .03 | .. | .. | .. | .. | .. |
| | 1.52 | 1.18 | .28 | 1.52 | .42 | 2.19 |

TABLE 3

Recalculated Analyses

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $SiO_2$ ..... | 59.21 | 57.85 | 60.74 | 63.34 | 64.59 | 65.86 |
| $TiO_2$ ..... | .99 | .98 | .80 | .82 | .70 | .60 |
| $Al_2O_3$ ..... | 20.31 | 20.81 | 18.54 | 16.51 | 16.60 | 15.73 |
| $Fe_2O_3$ ..... | 2.04 | 2.42 | 1.32 | 1.46 | 2.25 | 1.38 |
| FeO ..... | 4.99 | 5.41 | 5.60 | 5.83 | 3.93 | 4.19 |
| MnO ..... | .22 | .41 | .07 | .36 | .14 | .05 |
| MgO ..... | 1.88 | 1.78 | 2.42 | 2.22 | 2.05 | 1.51 |
| CaO ..... | .16 | .31 | 1.66 | 1.43 | 1.88 | 1.87 |
| BaO ..... | .10 | .10 | .... | .07 | .11 | .10 |
| $Na_2O$ ..... | 1.28 | 1.41 | 1.81 | 2.31 | 2.01 | 3.15 |
| $K_2O$ ..... | 4.56 | 4.34 | 4.03 | 3.21 | 3.83 | 3.52 |
| $H_2O^+$ ..... | 4.11 | 4.04 | 2.93 | 2.28 | 1.74 | 1.85 |
| $P_2O_5$ ..... | .15 | .14 | .08 | .16 | .17 | .19 |
| | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 |
| Sp. gr. ..... | 2.756 | 2.828 | 2.830 | 2.771 | 2.800 | 2.716 |

culated to represent compositions of 100 cc. rock.[2] The data are then compared with one another to determine additions and subtractions of material. Table 4 lists the results as applied to the Stavanger metamorphic series, while additions and subtractions between successive pairs of members of the series are given in table 5. It may be seen that no single constituent varies in the same manner throughout the series, but that any constituent may show gains in one or more pairs of rocks and losses in the next paired analyses.

Additions and subtractions are sometimes represented in percentages (Reynolds, 1943), and this has been done on an anhydrous basis in table 6. In the writer's opinion such representations are of little use and add no further information to the problem. Speculations regarding the composition of the pore fluid during the various metamorphic intensities can be drawn as well from table 5 as from table 6. Comparisons of the data of table 6 with compositions of magmas are probably unjustified and misleading. The time element is completely dis-

[2] In this method, and in others described later, sp. gr. data of rocks are used. It is doubtful whether the sp. gr. of a rock is generally determined with the same care and precision as is expended on its chemical analysis. A plea is here made for greater accuracy in the determination of this important physical constant of rocks.

TABLE 4

Compositions of 100 cc. Rock

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $SiO_2$ | 163.18 | 163.60 | 171.89 | 175.52 | 180.85 | 178.88 |
| $TiO_2$ | 2.73 | 2.77 | 2.26 | 2.27 | 1.96 | 1.63 |
| $Al_2O_3$ | 55.97 | 58.85 | 52.47 | 45.75 | 46.48 | 42.72 |
| $Fe_2O_3$ | 5.62 | 6.84 | 3.74 | 4.05 | 6.30 | 3.75 |
| FeO | 13.75 | 15.30 | 15.85 | 16.15 | 11.00 | 11.38 |
| MnO | .61 | 1.16 | .20 | 1.00 | .39 | .14 |
| MgO | 5.18 | 5.03 | 6.85 | 6.15 | 5.74 | 4.10 |
| CaO | .44 | .88 | 4.70 | 3.96 | 5.26 | 5.08 |
| BaO | .28 | .28 | .... | .19 | .31 | .27 |
| $Na_2O$ | 3.53 | 3.99 | 5.12 | 6.40 | 5.63 | 8.56 |
| $K_2O$ | 12.57 | 12.27 | 11.40 | 8.89 | 10.72 | 9.56 |
| $H_2O^+$ | 11.33 | 11.43 | 8.29 | 6.32 | 4.87 | 5.02 |
| $P_2O_5$ | .41 | .40 | .23 | .44 | .48 | .52 |
| | 275.60 | 282.80 | 283.00 | 277.09 | 279.99 | 271.61 |

TABLE 5

Additions and Subtractions*

| | 1-2 | | 2-3 | | 3-4 | | 4-5 | | 5-6 | |
|---|---|---|---|---|---|---|---|---|---|---|
| | + | − | + | − | + | − | + | − | + | − |
| $SiO_2$ | .42 | .. | 8.29 | .... | 3.63 | .... | 5.33 | ... | ... | 1.97 |
| $TiO_2$ | .04 | .. | .... | .51 | .01 | .... | .... | .31 | ... | .33 |
| $Al_2O_3$ | 2.88 | .. | .... | 6.38 | ... | 6.72 | .73 | ... | ... | 3.76 |
| $Fe_2O_3$ | 1.22 | .. | .... | 3.10 | .31 | .... | 2.25 | ... | ... | 2.55 |
| FeO | 1.55 | .. | .55 | .... | .30 | .... | .... | 5.15 | .38 | .... |
| MnO | .55 | .. | .... | .96 | .80 | .... | .... | .61 | ... | .25 |
| MgO | ... | .15 | 1.82 | .... | ... | .70 | .... | .41 | ... | 1.64 |
| CaO | .44 | .. | 3.82 | .... | ... | .74 | 1.30 | ... | ... | .18 |
| BaO | ... | .. | .... | .28 | .19 | .... | .12 | ... | ... | .04 |
| $Na_2O$ | .46 | .. | 1.13 | .... | 1.28 | .... | .... | .77 | 2.93 | .... |
| $K_2O$ | ... | .30 | .... | .87 | ... | 2.51 | 1.85 | ... | ... | 1.16 |
| $H_2O^+$ | .10 | .. | .... | 3.14 | ... | 1.97 | .... | 1.45 | .15 | .... |
| $P_2O_5$ | ... | .01 | .... | .17 | .21 | .... | .04 | ... | .04 | .... |
| | 7.66 | .46 | 15.61 | 15.41 | 6.73 | 12.64 | 11.60 | 8.70 | 3.50 | 11.88 |
| Less $H_2O^+$ | 7.56 | .46 | 15.61 | 12.27 | 6.73 | 10.67 | 11.60 | 7.25 | 3.35 | 11.88 |

*(+) and (−) symbols at top of columns indicate respectively additions and subtractions.

regarded in these speculations. Subtraction of, e.g., 5 per cent MgO does not mean that the resulting solution contained 5 per cent MgO, but that probably very dilute solutions in time removed 5 per cent MgO from the rock. Only the final results are observed in terms of rocks, and not in terms of pore fluids.

TABLE 6

Additions and Subtractions in Percentages
(on a Dry Basis)

| | 1-2 | | 2-3 | | 3-4 | | 4-5 | | 5-6 | |
|---|---|---|---|---|---|---|---|---|---|---|
| | + | − | + | − | + | − | + | − | + | − |
| $SiO_2$ | 5.56 | ..... | 53.11 | ..... | 53.93 | ..... | 45.95 | ..... | ..... | 16.58 |
| $TiO_2$ | .53 | ..... | ..... | 4.16 | .15 | ..... | ..... | 4.28 | ..... | 2.78 |
| $Al_2O_3$ | 38.10 | ..... | ..... | 52.00 | ..... | 62.98 | 6.29 | ..... | ..... | 31.66 |
| $Fe_2O_3$ | 16.14 | ..... | ..... | 25.26 | 4.61 | ..... | 19.40 | ..... | ..... | 21.46 |
| FeO | 20.49 | ..... | 3.52 | ..... | 4.46 | ..... | ..... | 71.03 | 11.34 | ..... |
| MnO | 7.28 | ..... | ..... | 7.82 | 11.89 | ..... | ..... | 8.41 | ..... | 2.10 |
| MgO | ..... | 32.61 | 11.66 | ..... | ..... | 6.56 | ..... | 5.66 | ..... | 13.80 |
| CaO | 5.82 | ..... | 24.47 | ..... | ..... | 6.94 | 11.21 | ..... | ..... | 1.52 |
| BaO | ..... | ..... | ..... | 2.28 | 2.82 | ..... | 1.03 | ..... | ..... | .34 |
| $Na_2O$ | 6.08 | ..... | 7.24 | ..... | 19.02 | ..... | ..... | 10.62 | 87.47 | ..... |
| $K_2O$ | ..... | 65.22 | ..... | 7.09 | ..... | 23.52 | 15.78 | ..... | ..... | 9.76 |
| $P_2O_5$ | ..... | 2.17 | ..... | 1.39 | 3.12 | ..... | .34 | ..... | 1.19 | ..... |
| | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 |

Joplin (1952) uses a slightly different method of calculation in which the rock exhibiting the lowest metamorphic grade is taken as standard and the percentages of the other analyses are multiplied with a factor reflecting the percentage increase or decrease in specific gravity as compared with that of the standard sample. The results of these calculations are listed in table 7. The present writer considers that there is no advantage in this manner of calculation instead of the more generally used method described above. The results of the calculations are ratio numbers and therefore less precise information is obtained by this method.

A third method has been seldom used (e.g., Ellis, 1947), probably because it requires more involved calculations and hence is more time-consuming. However, the use of a calculating machine largely alleviates the tediousness of the various computations. The atomic proportions are first calculated (table

TABLE 7

Equivalent Sp. Gr. Calculations

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $SiO_2$ | 59.21 | 59.36 | 62.37 | 63.68 | 65.62 | 64.91 |
| $TiO_2$ | .99 | 1.01 | .82 | .82 | .71 | .59 |
| $Al_2O_3$ | 20.31 | 21.35 | 19.04 | 16.60 | 16.87 | 15.50 |
| $Fe_2O_3$ | 2.04 | 2.48 | 1.36 | 1.47 | 2.29 | 1.36 |
| FeO | 4.99 | 5.55 | 5.75 | 5.86 | 3.99 | 4.13 |
| MnO | .22 | .42 | .07 | .36 | .14 | .05 |
| MgO | 1.88 | 1.83 | 2.49 | 2.23 | 2.08 | 1.49 |
| CaO | .16 | .32 | 1.70 | 1.44 | 1.91 | 1.84 |
| BaO | .10 | .10 | ..... | .07 | .11 | .10 |
| $Na_2O$ | 1.28 | 1.45 | 1.86 | 2.32 | 2.04 | 3.10 |
| $K_2O$ | 4.56 | 4.45 | 4.14 | 3.23 | 3.89 | 3.47 |
| $H_2O^+$ | 4.11 | 4.15 | 3.01 | 2.29 | 1.77 | 1.82 |
| $P_2O_5$ | .15 | .14 | .08 | .16 | .17 | .19 |
| | 100.00 | 102.61 | 102.69 | 100.53 | 101.59 | 98.55 |

8). $H_2O^+$ is reckoned as $2(OH)^{-1}$, the additional O being subtracted from the total oxygen proportion.[3] A conversion factor X is next found by dividing the specific gravity of the rock by $166.04\left(\frac{100}{\text{Avogadro number}}\right)$.[4] Multiplying the atomic proportions with the conversion factor finally gives the number of different atoms in 1 cc. of rock (table 10). The number of atoms added and subtracted in comparing rock pairs is given in table 11. Weights may be obtained from these figures by multiplying with the weight of the respective atoms. The sum of the weights of atoms should, of course, equal the specific gravity of the rock.

[3] The procedure followed here is slightly different from that normally used, in that $H_2O^+$ is reckoned as $2OH^{-1}$ and newer and more accurate physical constants are employed.

[4] Avogadro number and atomic weights from Handbook of Chemistry and Physics, 33rd ed., 1951-1952; ionic radii from Ahrens (1952).

TABLE 8

Atomic Proportions

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| Si | .9858 | .9632 | 1.0113 | 1.0546 | 1.0754 | 1.0966 |
| Ti | .0125 | .0123 | .0100 | .0103 | .0088 | .0075 |
| Al | .3984 | .4082 | .3638 | .3240 | .3256 | .3086 |
| $Fe^{+3}$ | .0256 | .0304 | .0166 | .0182 | .0282 | .0172 |
| $Fe^{+2}$ | .0694 | .0753 | .0779 | .0811 | .0547 | .0583 |
| Mn | .0032 | .0058 | .0010 | .0051 | .0020 | .0007 |
| Mg | .0466 | .0441 | .0600 | .0551 | .0508 | .0375 |
| Ca | .0029 | .0055 | .0296 | .0255 | .0335 | .0333 |
| Ba | .0007 | .0007 | .... | .0005 | .0007 | .0007 |
| Na | .0412 | .0454 | .0584 | .0746 | .0648 | .1016 |
| K | .0968 | .0922 | .0856 | .0682 | .0814 | .0748 |
| $OH^{-1}$ | .4568 | .4488 | .3256 | .2534 | .1934 | .2056 |
| P | .0022 | .0020 | .0012 | .0022 | .0024 | .0026 |
| $O^{-2}$ | 2.8299 | 2.8141 | 2.8567 | 2.8873 | 2.9199 | 2.9221 |
| Less O for OH | 2.6015 | 2.5897 | 2.6939 | 2.7606 | 2.8232 | 2.8193 |

TABLE 9

Physical Constants

Avogadro number: $6.0228 \times 10^{23}$

| Element | Ion. rad. (Å) | Ion. vol. 4.19 $r^3$ | At. wt. | Wt. 1 atom (gms) |
|---|---|---|---|---|
| $Si^{+4}$ | 0.42 | 0.31 | 28.06 | $46.60 \times 10^{-24}$ |
| $Ti^{+4}$ | 0.68 | 1.32 | 47.90 | 79.55 |
| $Al^{+3}$ | 0.51 | 0.56 | 26.97 | 44.80 |
| $Fe^{+3}$ | 0.64 | 1.10 | 55.85 | 92.76 |
| $Fe^{+2}$ | 0.74 | 1.70 | 55.85 | 92.76 |
| $Mn^{+2}$ | 0.80 | 2.15 | 54.93 | 91.21 |
| $Mg^{+2}$ | 0.66 | 1.20 | 24.32 | 40.39 |
| $Ca^{+2}$ | 0.99 | 4.07 | 40.08 | 66.56 |
| $Ba^{+2}$ | 1.34 | 10.08 | 137.36 | 228.10 |
| $Na^{+1}$ | 0.97 | 3.82 | 22.997 | 38.18 |
| $K^{+1}$ | 1.33 | 9.86 | 39.096 | 64.94 |
| $OH^{-1}$ | 1.40 | 11.48 | 17.008 | 28.24 |
| $P^{+5}$ | 0.35 | 0.18 | 30.98 | 51.44 |
| $O^{-2}$ | 1.40 | 11.48 | 16.000 | 26.57 |

TABLE 10

Number of Atoms in 1 cc. Rock

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| X | $1.66 \times 10^{22}$ | $1.70 \times 10^{22}$ | $1.70 \times 10^{22}$ | $1.67 \times 10^{22}$ | $1.69 \times 10^{22}$ | $1.64 \times 10^{22}$ |
| $N_o$ | $4.316 \times 10^{22}$ | $4.410 \times 10^{22}$ | $4.590 \times 10^{22}$ | $4.605 \times 10^{22}$ | $4.760 \times 10^{22}$ | $4.610 \times 10^{22}$ |
| $N_{si}$ | 1.635 | 1.640 | 1.723 | 1.759 | 1.813 | 1.793 |
| $N_{ti}$ | .021 | .021 | .017 | .017 | .015 | .012 |
| $N_{al}$ | .661 | .695 | .620 | .540 | .549 | .505 |
| $N_{fe^{+3}}$ | .042 | .052 | .028 | .030 | .048 | .028 |
| $N_{fe^{+2}}$ | .115 | .128 | .133 | .135 | .092 | .095 |
| $N_{mn}$ | .005 | .010 | .002 | .009 | .003 | .001 |
| $N_{mg}$ | .077 | .075 | .102 | .092 | .086 | .061 |
| $N_{ca}$ | .005 | .009 | .050 | .043 | .056 | .054 |
| $N_{ba}$ | .001 | .001 | ... | .001 | .001 | .001 |
| $N_{na}$ | .068 | .077 | .100 | .124 | .109 | .166 |
| $N_k$ | .161 | .157 | .146 | .114 | .137 | .122 |
| $N_{oh}$ | .758 | .764 | .555 | .423 | .326 | .336 |
| $N_p$ | .004 | .003 | .002 | .004 | .004 | .004 |
| | 7.869 | 8.042 | 8.068 | 7.896 | 7.999 | 7.788 |

TABLE 11

Additions and Subtractions

(All Numbers x $10^{22}$)

| | 1-2 | | 2-3 | | 3-4 | | 4-5 | | 5-6 | |
|---|---|---|---|---|---|---|---|---|---|---|
| | + | − | + | − | + | − | + | − | + | − |
| O | .094 | ... | .180 | ... | .015 | ... | .155 | ... | ... | .150 |
| Si | .005 | ... | .083 | ... | .036 | ... | .054 | ... | ... | .020 |
| Ti | ... | ... | ... | .004 | ... | ... | ... | .002 | ... | .003 |
| Al | .034 | ... | ... | .075 | ... | .080 | .009 | ... | ... | .044 |
| $Fe^{+3}$ | .010 | ... | ... | .024 | .002 | ... | .018 | ... | ... | .020 |
| $Fe^{+2}$ | .013 | ... | .005 | ... | .002 | ... | ... | .043 | .003 | ... |
| Mn | .005 | ... | ... | .008 | .007 | ... | ... | .006 | ... | .002 |
| Mg | ... | .002 | .027 | ... | ... | .010 | ... | .006 | ... | .025 |
| Ca | .004 | ... | .041 | ... | ... | .007 | .013 | ... | ... | .002 |
| Ba | ... | ... | ... | .001 | .001 | ... | ... | ... | ... | ... |
| Na | .009 | ... | .023 | ... | .024 | ... | ... | .015 | .057 | ... |
| K | ... | .004 | ... | .011 | ... | .032 | .023 | ... | ... | .015 |
| OH | .006 | ... | ... | .209 | ... | .132 | ... | .097 | .010 | ... |
| P | ... | .001 | ... | .001 | .002 | ... | ... | ... | ... | ... |
| | .180 | .007 | .359 | .333 | .089 | .261 | .272 | .169 | .070 | .281 |
| Vals. | .011 | .013 | .141 | .138 | .174 | .174 | .036 | .036 | .053 | .057 |

In table 12 the ion volumes have been listed, as obtained by multiplying the number of atoms with the corresponding ionic volume (table 9). Summation of the ion volumes gives a measure of the packing of the rock comparable with the packing index concept $\left(\frac{\text{ion volume x 10}}{\text{volume rock}}\right)$ introduced by Fairbairn (1943). When comparing packing percentages with packing indices given by Fairbairn, it should be noted that different values are used for ionic volumes. This is especially significant in the case of oxygen (11.48 as against 9.64 cubic Å). Again no regular trend can be discerned in the packing variation of the six rocks.

TABLE 12

Ion Volumes in 1 cc. Rock

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $V_o$ ..... | .495 cc | .506 cc | .527 cc | .529 cc | .546 cc | .529 cc |
| $V_{si}$ ..... | .005 | .005 | .005 | .006 | .006 | .006 |
| $V_{ti}$ ..... | .000 | .000 | .000 | .000 | .000 | .000 |
| $V_{al}$ ..... | .004 | .004 | .004 | .003 | .003 | .003 |
| $V_{fe^{+3}}$ ..... | .001 | .001 | .000 | .000 | .001 | .000 |
| $V_{fe^{+2}}$ ..... | .002 | .002 | .002 | .002 | .002 | .002 |
| $V_{mn}$ ..... | .000 | .000 | .000 | .000 | .000 | .000 |
| $V_{mg}$ ..... | .001 | .001 | .001 | .001 | .001 | .001 |
| $V_{ca}$ ..... | .000 | .000 | .002 | .002 | .002 | .002 |
| $V_{ba}$ ..... | .000 | .000 | .000 | .000 | .000 | .000 |
| $V_{na}$ ..... | .003 | .003 | .004 | .005 | .004 | .006 |
| $V_k$ ..... | .016 | .016 | .014 | .011 | .014 | .012 |
| $V_{oh}$ ..... | .087 | .088 | .064 | .049 | .037 | .039 |
| $V_p$ ..... | .000 | .000 | .000 | .000 | .000 | .000 |
| | .614 cc | .626 cc | .623 cc | .608 cc | .616 cc | .600 cc |
| Packing (%) | 61.4 | 62.6 | 62.3 | 60.8 | 61.6 | 60.0 |

CALCULATIONS BASED ON A "UNIT CELL" OF 160 O ATOMS

Barth (1948) introduced a new method of calculation based on no less than three premises: (1) that in metasomatic processes the volume remains constant, (2) that the number of oxygen atoms also remains constant, and (3) that rocks generally contain 100 cations to 160 oxygen atoms. Thus the analyses are recalculated to a unit cell of 160 O atoms, for which Barth uses the total available oxygen. The results of these calculations for the six Stavanger rocks are shown in table 13, while pairs of rocks are compared in table 14.

TABLE 13

Analyses Recalculated to 160 O Unit Cell

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| Si | 51.57 | 50.72 | 53.59 | 55.98 | 57.04 | 58.00 |
| Ti | .65 | .65 | .53 | .55 | .47 | .40 |
| Al | 20.84 | 21.49 | 19.28 | 17.20 | 17.27 | 16.32 |
| $Fe^{+3}$ | 1.34 | 1.60 | .88 | .97 | 1.50 | .91 |
| $Fe^{+2}$ | 3.63 | 3.97 | 4.13 | 4.31 | 2.90 | 3.08 |
| Mn | .17 | .31 | .05 | .27 | .11 | .04 |
| Mg | 2.44 | 2.32 | 3.18 | 2.93 | 2.69 | 1.98 |
| Ca | .15 | .29 | 1.57 | 1.35 | 1.78 | 1.76 |
| Ba | .04 | .04 | ..... | .03 | .04 | .04 |
| Na | 2.16 | 2.39 | 3.09 | 3.96 | 3.44 | 5.37 |
| K | 5.06 | 4.86 | 4.54 | 3.62 | 4.32 | 3.96 |
| H | 23.90 | 23.63 | 17.25 | 13.45 | 10.26 | 10.88 |
| P | .12 | .11 | .06 | .12 | .13 | .14 |
| | 112.07 | 112.38 | 108.15 | 104.74 | 101.95 | 102.74 |

As shown in the last table, the total charge of the ions added roughly equals that of the subtracted ions; thus electrostatic equilibrium is maintained during metasomatism. The same approximate balance of valences may be observed in the addition and subtraction of atoms when comparing their numbers in 1 cc. rock (table 11). Comparison of tables 5, 11, and 14 shows discrepancies in the addition or subtraction of $Si^{+4}$ and $H^{+1}$.

TABLE 14

Additions and Subtractions

| | 1-2 | | 2-3 | | 3-4 | | 4-5 | | 5-6 | |
|---|---|---|---|---|---|---|---|---|---|---|
| | + | − | + | − | + | − | + | − | + | − |
| Si | ... | .85 | 2.87 | ... | 2.39 | ... | 1.06 | ... | .96 | ... |
| Ti | ... | ... | ... | .12 | .02 | ... | ... | .08 | ... | .07 |
| Al | .65 | ... | ... | 2.21 | ... | 2.08 | .07 | ... | ... | .95 |
| $Fe^{+3}$ | .26 | ... | ... | .72 | .09 | ... | .53 | ... | ... | .59 |
| $Fe^{+2}$ | .34 | ... | .16 | ... | .22 | ... | ... | 1.41 | .18 | ... |
| Mn | .14 | ... | ... | .26 | .18 | ... | ... | .16 | ... | .07 |
| Mg | ... | .12 | .86 | ... | ... | .25 | ... | .24 | ... | .71 |
| Ca | .14 | ... | 1.28 | ... | ... | .22 | .43 | ... | ... | .02 |
| Ba | ... | ... | ... | .04 | .03 | ... | .01 | ... | ... | ... |
| Na | .23 | ... | .70 | ... | .87 | ... | ... | .52 | 1.93 | ... |
| K | ... | .20 | ... | .32 | ... | .92 | .70 | ... | ... | .36 |
| H | ... | .27 | ... | 6.38 | ... | 3.80 | ... | 3.19 | .62 | ... |
| P | ... | .01 | ... | .05 | .06 | ... | .01 | ... | .01 | ... |
| | 1.76 | 1.45 | 5.87 | 10.10 | 3.86 | 5.27 | 2.81 | 5.60 | 3.70 | 2.77 |
| Vals. | 4.20 | 4.16 | 16.78 | 16.82 | 11.94 | 11.90 | 7.67 | 7.65 | 6.80 | 6.86 |

Inspection of table 10 shows that Barth's second premise is not correct when taken in conjunction with the first; constant volume does not necessarily mean a constant number of oxygen atoms. On these and other grounds the Barth method of calculation has been criticized severely by Rosenqvist (1949) and Brajnikov (1949).

Goldschmidt (1928) has emphasized the role of oxygen in rocks; the lithosphere is essentially an "oxysphere." Bragg (1937, p. 31) states that "oxygen is the prominent union in most minerals. It is so large that the packing together of the ions in the structures is mainly a packing of oxygen ions, with the metallic cations in the interstices." Read (1948, p. 180) suggests that in metasomatic processes such as metamorphism and granitization "the large oxygen atoms of the silicate lattice remain more or less in position whilst the small migrating atoms drift through." It is on these authoritative statements that Barth bases his assertion that the volume of a rock is essentially the volume of the oxygen atoms in the rock, and that therefore constant volume means a constant number of oxygen atoms. This argument ignores the fact that a considerable proportion of any mineral or rock consists of empty space, and that due to different ionic packing, the proportion of empty space is likely to vary from one mineral to another, or from one rock to another.

The present writer, while recognizing the predominant role played by oxygen in silicate structures, is not convinced that oxygen remains at a constant level during metasomatism. The large size of the oxygen ion does not provide a valid argument, because the potash ion is nearly as large and yet potash apparently migrates with great ease. In the rearrangement of the $(Si, Al)O_4$ tetrahedra, resulting in the formation of new minerals, oxygen is liberated or introduced, presumably combining in or produced from $OH^{-1}$ ions present in the ubiquitous aqueous fluid phase. Much mineralogical evidence points toward the high stability of the $(Si, Al)O_4$ tetrahedral structure and the high energy barrier to be overcome in destroying this structure. Apparently even in magmas $SiO_4$ tetrahedra are present. It appears therefore more logical to assume that the $(Si, Al)O_4$ building blocks of the silicate structure are merely rearranged to a different pattern and remain at an approximately constant

number in metasomatism, than that the number of oxygen atoms remains constant.

An illustration of this view is provided by the coronas developed in metamorphosed rocks consisting of magnesian olivine and calcic plagioclase only. Magnesian orthopyroxene generally occurs as the first rim around olivine crystals and is formed at the expense of that mineral. The orthodox equation for the reaction is

$$(Mg,Fe)_2SiO_4 + SiO_2 \rightarrow 2\ (Mg,Fe)SiO_3.$$

The source of the silica required for this reaction is a mystery in the present case. Shand (1945) has also expressed dissatisfaction with current equations explaining the formation of coronas. In the opinion of the present writer the formation of orthopyroxene from olivine results in this case from a rearrangement of the independent tetrahedra sharing no oxygen in olivine, to a pattern consisting of chains of tetrahedra, each sharing two oxygen atoms, in orthopyroxene. Together with this rearrangement of the fundamental building blocks $Mg^{+2}$ and $O^{-2}$ migrated outwards, the oxygen combining in $OH^{-1}$ ions. $SiO_4$ tetrahedra were neither added nor subtracted in the process. If this is written at all in the form of a chemical equation, some such representation as the following would seem to be preferable to the orthodox equation

$$(Mg,Fe)_2SiO_4 + H_2O \rightarrow (Mg,Fe)SiO_3 + (Mg,Fe)^{+2} + 2OH^{-1}.$$

### CONSTANT VOLUME IN METAMORPHISM AND GRANITIZATION

The law of equal volume has dominated petrological thought concerning metasomatic processes for a considerable length of time. It is based on such observations as the perfect preservation of fossils in metamorphic minerals (Bucher, 1953) and the pseudomorphous replacement of one mineral by another, both without sign of cracks or open spaces to indicate volume changes. A good case can therefore be made for constant volume relations in detail, but it remains uncertain whether its extension to rocks showing evidence of regional metamorphism and granitization is fully justified.

Acceptance of constant volume relations has led to acceptance of external rather than internal forces to explain the

intricate foldings so commonly observed in metamorphic terrains. Thus the remarkable thinning of highly metamorphosed pelitic beds is attributed to tight squeezing, and the up-doming of migmatized sediments is ascribed to the upward pressures of ascending magmas which remain hidden from direct observation. Whatever changes in volume did occur were the consequence of externally applied tension or compression, rather than the result of recrystallization and changes in composition within the rocks themselves. The logical conclusion is that regional metamorphism is the outcome of orogenesis by such external forces, rather than that the foldings observed in plutonic terrains result from metamorphism, an interesting viewpoint advanced by Read (1952).

Not all authorities have thus widely applied the law of equal volume. Van Hise (1904) assumed that metamorphism is normally attended by volume changes which can be calculated by the application of Lepsius' volume law (mol. vol. $= \frac{\text{mol. weight}}{\text{sp. gr.}}$). Irving (1911, pp. 664-665) argued that replacement is frequently accompanied by a decrease in volume, as shown by vugs in the new rock, but significantly states: "It is . . . impossible to calculate this change exactly from analyses, unless some one element has persisted unchanged in the process . . . which does not often happen. Whatever the theoretically calculated change, the observed facts show that in nearly all siliceous replacements and in the greater number of metallic replacements, there has been an appreciable and often marked decrease in the volume of the rock." More recently Eskola (1948) suggested that granitization of sediments causes an increase in volume whereby the rocks are arched up in mantled gneiss domes. Perrin (1950) employed the calculation methods of Brajnikov (1949) and Barth (1948) in an attempt to determine volume changes resulting from granitization processes, using analyses given by Grout (1937) and Anderson (1937) as examples. Those favoring volume changes in metamorphic and granitization metasomatism are distinctly in the minority today, and are by no means agreed that a volume increase or decrease is the outcome of such processes.

The volume of a rock really incorporates several variable components, the most significant of which are: (1) the ion

volumes, (2) the theoretical empty space resulting from ideal packing of the ions in different minerals, (3) the disorder empty space, and (4) the pore space which also includes the minute capillary and sub-capillary openings normally not measured in porosity determinations. Metasomatism resulting from regional metamorphism means essentially the invasion of a rock by an active, aqueous fluid of low viscosity, consequent to a rise in temperature level. This fluid must gain access to every part of the rock, hence it is reasonable to suppose that a large number of openings were available in the rock at the time of replacement. Many of these minute cracks probably were healed at a later stage in a similar fashion as the cracks in quartz from Washington (D. C.) were healed, leaving trails of fluid inclusions as only evidence of their erstwhile existence (Tuttle, 1949). Metasomatism probably is attended by an initial increase in both pore space and disorder empty space, which may be augmented by an increase, or offset by a decrease in ion volume and theoretical empty space. The assumption that throughout the process increases of some of the volume components are always exactly balanced by decreases in the other components, so that the bulk volume remains the same, is scarcely to be credited.

It appears impossible to calculate with accuracy the actual change in volume which might occur upon the metasomatic transformation of any given rock. On the basis of constant oxygen or constant number of $(Si,Al)O_4$ tetrahedra we may calculate apparent volume changes, but the results are representative of only two of the four volume components, and completely ignore volume changes resulting from differences in disorder empty space and pore space. It is also likely that during lengthy processes such as metamorphism, metasomatism, or granitization, there are several changes in volume which may in part counteract one another. Field studies may indicate such changes in volume, and the present writer attaches much greater significance to properly presented field evidence of volume changes than to hypothetical volume changes calculated, by whatever method, from chemical analyses. However, rigid adherence to the law of equal volume inevitably results in translation of the field evidence in terms of externally applied forces. Even when an unbiased attitude is preserved, field relations may yield apparently conflicting results. For instance, an initial decrease

in volume may be followed by a later increase in volume, or vice versa, during the same metamorphic cycle. Yet the writer believes that the problem of volume changes during metamorphism is probably attacked best by detailed studies of field relations. Evidence from orogenic belts of alternating periods of compression and tension might perhaps be interpreted as due to volume changes resulting from metamorphism and granitization.

### CALCULATIONS BASED ON A CONSTANT NUMBER OF (SI,AL)$O_4$ TETRAHEDRA

Modern mineralogical and petrological theory evokes a picture in which changes in composition of rocks and changes of one mineral to another are the result of a rearrangement in the configuration of the fundamental building blocks of (Si,Al)$O_4$ tetrahedra, accompanied by removal, substitution, or introduction of the interstitial cations. The most probable conclusion is that in these changes the number of (Si,Al)$O_4$ tetrahedra remains the same, although there may be partial substitution of $Si^{+4}$ by $Al^{+3}$ in the tetrahedra. The problem is therefore to find a method which allows for the calculation of the approximate number of tetrahedra in a rock. If the mode and chemical analyses of the constituent minerals are available in addition to the chemical analysis of the rock, $Si^{+4}$-$Al^{+3}$ substitution and the number of (Si,Al)$O_4$ tetrahedra can be calculated with accuracy. In the absence of chemical analyses of the constituent minerals, the geologic literature can be consulted for reliable mineral analyses from comparable rocks, in order to determine $Si^{+4}$-$Al^{+3}$ substitution. If, as in the present case, the modes of the rocks are also lacking, a less accurate method of calculation must be used. In the C.I.P.W. calculation of the norm, the alkalies and calc-alkalies, after allotment for certain minor constituents, are used up in the formation of feldspars, the excess alumina, if any, being assumed to exist as corundum. A similar procedure is followed in table 15, based on the atomic proportions listed in table 8. Part of the Ca is allotted for apatite in the ratio P : Ca = 2 : 3, while the remaining (Ca+Ba) is allotted for anorthite in the ratio (Ca+Ba) : Al = 1 : 2. The combined alkalies are allotted for albite and orthoclase in the ratio (Na+K) : Al = 1 : 1. Aluminum used for feldspar is assumed to represent the total aluminum substituting for silicon in tetrahedral posi-

TABLE 15

Calculation of $(Si,Al)O_4$ Tetrahedra Number

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| Ca | .0029 | .0055 | .0296 | .0255 | .0335 | .0333 |
| P | .0022 | .0020 | .0012 | .0022 | .0024 | .0026 |
| Ca for Ap. | .0029 | .0030 | .0018 | .0033 | .0036 | .0039 |
| Rest Ca | .0000 | .0025 | .0278 | .0222 | .0299 | .0294 |
| Ba | .0007 | .0007 | .0000 | .0005 | .0007 | .0007 |
| Al for An. | .0014 | .0064 | .0556 | .0454 | .0612 | .0602 |
| Na | .0412 | .0454 | .0584 | .0746 | .0648 | .1016 |
| K | .0968 | .0922 | .0856 | .0682 | .0814 | .0748 |
| Al for Ab.+Or. | .1380 | .1376 | .1440 | .1428 | .1462 | .1764 |
| Al repl. Si | .1394 | .1440 | .1996 | .1882 | .2074 | .2366 |
| Si | .9858 | .9632 | 1.0113 | 1.0546 | 1.0754 | 1.0966 |
| Proportion $(Si,Al)O_4$ | 1.1252 | 1.1072 | 1.2109 | 1.2428 | 1.2828 | 1.3332 |
| Factor X | $1.66 \times 10^{22}$ | $1.70 \times 10^{22}$ | $1.70 \times 10^{22}$ | $1.67 \times 10^{22}$ | $1.69 \times 10^{22}$ | $1.64 \times 10^{22}$ |
| Number of $(Si,Al)O_4$ in 1 cc | $1.867 \times 10^{22}$ | $1.886 \times 10^{22}$ | $2.063 \times 10^{22}$ | $2.073 \times 10^{22}$ | $2.180 \times 10^{22}$ | $2.180 \times 10^{22}$ |

TABLE 16

Number of Atoms to Constant $(Si,Al)O_4$ Tetrahedra Number
$2.180 \times 10^{22}$

| | 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|---|
| $N_o$ | $5.040 \times 10^{22}$ | $5.098 \times 10^{22}$ | $4.849 \times 10^{22}$ | $4.842 \times 10^{22}$ | $4.760 \times 10^{22}$ | $4.610 \times 10^{22}$ |
| $N_{si}$ | 1.909 | 1.896 | 1.820 | 1.850 | 1.813 | 1.793 |
| $N_{ti}$ | .025 | .024 | .018 | .018 | .015 | .012 |
| $N_{al}$ | .772 { .271 / .501 } | .803 { .284 / .519 } | .655 { .360 / .295 } | .568 { .330 / .238 } | .549 { .367 / .182 } | .505 { .387 / .118 } |
| $N_{fe^{+3}}$ | .049 | .060 | .030 | .032 | .048 | .028 |
| $N_{fe^{+2}}$ | .134 | .148 | .141 | .142 | .092 | .095 |
| $N_{mn}$ | .006 | .012 | .002 | .009 | .003 | .001 |
| $N_{mg}$ | .090 | .087 | .108 | .097 | .086 | .061 |
| $N_{ca}$ | .006 | .010 | .053 | .045 | .056 | .054 |
| $N_{ba}$ | .001 | .001 | .... | .001 | .001 | .001 |
| $N_{na}$ | .079 | .089 | .106 | .130 | .109 | .166 |
| $N_k$ | .188 | .181 | .154 | .120 | .137 | .122 |
| $N_{oh}$ | .885 | .883 | .586 | .445 | .326 | .336 |
| $N_p$ | .005 | .003 | .002 | .004 | .004 | .004 |
| | $9.189 \times 10^{22}$ | $9.295 \times 10^{22}$ | ——— | ——— | ——— | ——— |
| | ——— | ——— | $8.524 \times 10^{22}$ | $8.303 \times 10^{22}$ | $7.999 \times 10^{22}$ | $7.788 \times 10^{22}$ |

TABLE 17

Additions and Subtractions
(All Numbers x $10^{22}$)

| | 1-2 | | 2-3 | | 3-4 | | 4-5 | | 5-6 | |
|---|---|---|---|---|---|---|---|---|---|---|
| | + | − | + | − | + | − | + | − | + | − |
| O | .058 | ... | ... | .249 | ... | .007 | ... | .082 | ... | .150 |
| Si | ... | .013 | ... | .076 | .030 | ... | ... | .037 | ... | .020 |
| Ti | ... | .001 | ... | .006 | ... | ... | ... | .003 | ... | .003 |
| Al | .013 | ... | .076 | ... | ... | .030 | .037 | ... | .020 | ... |
| | .018 | ... | ... | .224 | ... | .057 | ... | .056 | ... | .064 |
| $Fe^{+3}$ | .011 | ... | ... | .030 | .002 | ... | .016 | ... | ... | .020 |
| $Fe^{+2}$ | .014 | ... | ... | .007 | .001 | ... | ... | .050 | .003 | ... |
| Mn | .006 | ... | ... | .010 | .007 | ... | ... | .006 | ... | .002 |
| Mg | ... | .003 | .021 | ... | ... | .011 | ... | .011 | ... | .025 |
| Ca | .004 | ... | .043 | ... | ... | .008 | .011 | ... | ... | .002 |
| Ba | ... | ... | ... | .001 | .001 | ... | ... | ... | ... | ... |
| Na | .010 | ... | .017 | ... | .024 | ... | ... | .021 | .057 | ... |
| K | ... | .007 | ... | .027 | ... | .034 | .017 | ... | ... | .015 |
| OH | ... | .002 | ... | .297 | ... | .141 | ... | .119 | .010 | ... |
| P | ... | .002 | ... | .001 | .002 | ... | ... | ... | ... | ... |
| | .134 | .028 | .081 | .852 | .067 | .288 | .044 | .348 | .070 | .281 |
| Vals. | .068 | .077 | .145 | .135 | .178 | .178 | .087 | .089 | .053 | .057 |

tion, the remaining aluminum being interstitial to the tetrahedra. Thus the proportion of $(Si,Al)O_4$ tetrahedra is represented by the atomic proportion of total Si plus Al used in feldspar. From this proportion the number of tetrahedra in 1 cc. of rock is found by multiplying with the conversion factor X (table 10). Table 16 lists the number of atoms, assuming a constant number of $(Si,Al)O_4$ tetrahedra of $2.180 \times 10^{22}$, the number found for the highest grade of the series.[5] Table 17 shows additions and subtractions of atoms obtained by comparing pairs of rocks of table 16. As may be expected, table 17 differs materially from table 11. Weights may again be calculated from the number of atoms listed in table 17, by multiplying with the weight of the corresponding atom (table 9).

Several significant features of the changes of composition in the six rocks can be deduced from table 16. First, it is clear that there is a general tendency toward decrease in oxygen; total oxygen, oxygen not bound in $OH^{-1}$, as well as oxygen bound in $OH^{-1}$. This means that in this rock series there is an increase in the number of oxygen atoms shared between $(Si,Al)O_4$ tetrahedra. Second, the number of silicon atoms decreases in the rock series, in spite of the number of $(Si,Al)O_4$ tetrahedra being kept constant, and in spite of the fact that the percentage silica of the original analyses shows a marked increase. The decrease in silicon at constant number of tetrahedra is explained by the progressive increase in $Si^{+4}$-$Al^{+3}$ substitution. There is a progressive decrease in total aluminum, compounded of an increase in the number of Al atoms substituting for Si in the tetrahedral structure, and a decrease in the number of Al atoms in 6-coordination. Third, the number of cations interstitial to the $(Si,Al)O_4$ tetrahedra shows a progressive decrease in the rock series. The fact that there is a progressive decrease in both total oxygen and in the number of interstitial cations explains why there should be an increase in the percentage of silica in the original analyses, although the number of silicon atoms decreases. The rocks have been relatively enriched in silica, yet silicon atoms were not added but *removed* in the process.

We may recalculate the analyses also to a hypothetical unit cell consisting of a certain number of $(Si,Al)O_4$ tetrahedra.

[5] It is purely a matter of choice to which number of $(Si, Al)O_4$ tetrahedra the data are recalculated.

This considerably simplifies the computations, but the results are again qualitative rather than quantitative, and may even be misleading. Barth (1948) used a similar procedure in his assumption of constant number of oxygen atoms, and chose a unit cell of 160 O atoms, with the further assumption that rocks generally contain 100 cations to 160 O atoms. It is easily calculated from table 10 that in the present case the number of oxygen atoms per 100 cations varies from 142 for analysis 2, to 157 for analysis 5. A unit cell of a certain number of $(Si,Al)O_4$ tetrahedra would be entirely artificial and would have no constant relation to the number of included cations or oxygen atoms. Hence the present writer cannot recommend such a procedure.

In replacement ore bodies $(Si,Al)O_4$ tetrahedra have evidently been removed. Similarly, in weathering the tetrahedral building blocks of silicates may be partly broken down and removed. However, these processes are perhaps not the same as the metasomatic changes in chemical compositions of rocks which accompany metamorphism and granitization. Weathering proceeds at lower energy levels than apply to metamorphism and granitization. Moreover, rock-reconstitutions in metamorphism are usually effected in response to a gradually rising energy level, while replacement ore bodies are believed to have been formed in an environment of slowly diminishing temperatures. The assumption of a constant number of $(Si,Al)O_4$ tetrahedra clearly cannot be applied to *all* metasomatic processes, but it is thought to be valid when restricted to metamorphism, granitization, and similar replacements of silicates by other silicates in the absence of actual melting.

Metamorphism and granitization are here regarded as essentially the result of increasing temperatures. This is attended by an initial rapid increase in pore spaces, affording passages for invading, active fluids, the media of replacement. Changes of volume necessarily occur as the outcome of increase in pore space and disorder empty space, recrystallization, and ionic substitution, addition, or subtraction, but such changes can only be predicted qualitatively from field relations, and probably vary even during the different stages of the process. Volume changes result in the deformation of the rocks, rendered plastic by the existing higher temperatures and permeation by the aqueous fluids. Thus, the intricate foldings often observed in

metamorphic terrains are believed to be, at least in part, the outcome of these volume changes which in turn result from the rise in temperature. This last factor initiates metamorphism and granite formation.

## CONCLUSIONS

Doubts are expressed in this paper that either constant volume relations or a constant number of oxygen atoms invariably prevail in regional metamorphism and granitization. These assumptions form the basis of all current methods of calculating changes in bulk composition of rocks in their transformation from one metamorphic grade or phase of granitization to another. Instead, it is suggested that the number of $(Si,Al)O_4$ tetrahedra remains constant in these processes, which are further marked by partial substitution of $Si^{+4}$ by $Al^{+3}$, rearrangement of the tetrahedral building blocks of silicate structures to different patterns, and substitution, introduction, or removal of the interstitial ions. A method is suggested whereby changes in bulk composition of rocks during these processes may be calculated. Volume changes which attended the transformation of the rocks may be calculated on the same basis, but the results are not believed to have much significance.

In the writer's view, metasomatic transformations of rocks as a result of regional metamorphism, are heralded by an increase in temperature and the soaking of the rocks by an active, aqueous fluid which permeates the rocks through passages opened at least partly in response to rising temperature. Changes in volume of the rocks may be responsible for their deformation, and in the case of granitized rocks, for their eventually assuming intrusive relations towards the associated rocks. Thus a great deal of the foldings exhibited by plutonic terrains may be the consequence of metamorphism and granitization, instead of their being the outcome of external forces which also resulted in metamorphism and granitization.

## ACKNOWLEDGMENTS

The writer has benefited greatly from many discussions with his students and with geologists interested in the problems of regional metamorphism. Drs. Ch. H. Behre, W. H. Bucher, A. F. Buddington, H. H. Hess, and H. Kuno critically read the first draft and made many valuable suggestions. The views

expressed in this paper are entirely the writer's, but he is very appreciative of the kind interest shown and helpful comments made by these fellow geologists.

### References

Ahrens, L. H., 1952. The use of ionization potentials. Part 1. Ionic radii of the elements: Geochim. Cosmochim. Acta, vol. 2, pp. 155-169.

Anderson, G. H., 1937. Granitization, albitization and related phenomena in the Inyo Range of California-Nevada: Geol. Soc. America Bull., vol. 48, pp. 1-74.

Barth, T. F. W., 1948. Oxygen in rocks: A basis for petrographic calculations: Jour. Geology, vol. 56, pp. 50-60.

Bragg, W. L., 1937. Atomic structure of minerals, Cornell University Press, Ithaca, N. Y.

Brajnikov, B., 1949. Les rapports numériques et volumiques de l'oxygêne dans les minéraux et les roches: Instituto de Tecnologia Industrial, Minas Gerais, Bol. 8.

Bucher, W. H., 1953. Fossils in metamorphic rocks: A review: Geol. Soc. America Bull., vol. 64, pp. 275-300.

Chao, E. C. T., 1951. Granitization and basification by diffusion: Norsk geol. tidsskr., vol. 29, pp. 84-107.

Ellis, J., 1947. The Marievale granophyre—a metamorphic rock: Geol. Soc. South Africa Trans., vol. 50, pp. 121-160.

Eskola, P. E., 1948. The problem of mantled gneiss domes: Geol. Soc. London Quart. Jour., vol. 104, pp. 461-476.

Faber, H., 1941. On the salt solutions in microscopic cavities in granites: Danmarks geol. Undersögelse, vol. 2, no. 67.

Fairbairn, H. W., 1943. Packing in ionic minerals: Geol. Soc. America Bull., vol. 54, pp. 1305-1374.

Goldschmidt, V. M., 1920. Die Injektionsmetamorphose im Stavanger-Gebiete. Geologisch-petrographische Studien im Hochgebirge des südlichen Norwegens. V: Norske vidensk.-akad. Skr., Mat.-Nat. Kl., no. 10.

———, 1928. Ueber die Raumerfüllung der Atome (Ionen) in Kristallen und über das Wesen der Lithosphäre: Neues Jahrb., Beilage-Band 57, Abt. A, pp. 1119-1130.

Grout, F. F., 1937. Criteria of inclusions in plutonic rocks: Geol. Soc. America Bull., vol. 48, pp. 1521-1572.

Irving, J. D., 1911. Replacement ore-bodies and the criteria for their recognition: Econ. Geology, vol. 6, pp. 527-561, 619-669.

Joplin, G. A., 1952. The granitization process and its limitations as exemplified in certain parts of New South Wales: Geol. Mag., vol. 89, pp. 25-38.

Lindgren, W., 1933. Mineral deposits, 4th ed., McGraw-Hill Book Company, New York.

Perrin, P., 1950. L'oxygêne en les calculs pétrographiques, une discussion: Jour. Geology, vol. 58, pp. 163-168.

Read, H. H., 1948. A commentary on place in plutonism: Geol. Soc. London Quart. Jour., vol. 104, pp. 155-205 (presidential address).

———, 1952. Metamorphism and granitization: Geol. Soc. South Africa Trans., vol. 54, suppl.

Reynolds, D. L., 1943. The south-western end of the Newry Igneous Complex. A contribution towards the petrogenesis of the granodiorites: Geol. Soc. London Quart. Jour., vol. 99, pp. 205-246.

———, 1946. The sequence of geochemical changes leading to granitization: Geol. Soc. London Quart. Jour., vol. 102, pp. 389-446.

Rosenqvist, I. Th., 1949. The distribution of oxygen in the lithosphere and oxygen in rocks; a basis for petrographic calculations. A discussion: Jour. Geology, vol. 57, pp. 420-423.

Shand, S. J., 1945. Coronas and coronites: Geol. Soc. America Bull., vol. 56, pp. 247-266.

Tuttle, O. F., 1949. Structural petrology of planes of liquid inclusions: Jour. Geology, vol. 57, pp. 331-356.

Van Hise, C. R., 1904. A treatise on metamorphism: U. S. Geol. Survey Mon., 47.

Department of Geology
Columbia University
New York, N. Y.

[American Journal of Science, Vol. 251, July 1953, Pp. 505-542]

# POSTGLACIAL FORESTS IN THE YUKON TERRITORY AND ALASKA*

HENRY P. HANSEN

ABSTRACT. Pollen analysis of peat sections from 74 sites of organic sedimentation in the Yukon Territory and Alaska reveals a record of the forests that have existed within range of pollen dispersal for all or most of the time since this region was freed of the last glacier. Sections along the Alaska Highway in the Yukon indicate that lodgepole pine (*Pinus contorta latifolia*) has been abundant during the time represented and predominant over spruce throughout most of the sections as far as milepost 931, about 15 miles northwest of Whitehorse. The present range of lodgepole pine does not extend far beyond this point, indicating that its postglacial range has not differed significantly from its present range. The early appearance of pine pollen in these sections substantiates theories and evidence that pine persisted in unglaciated areas in west-central Yukon during the Wisconsin glacial age.

From milepost 931 to the Yukon-Alaska border, and throughout interior Alaska, spruce has been the predominant forest tree, and in most sections it is recorded to nearly 100 per cent. In the Fairbanks and Anchorage regions white birch (*Betula papyrifera* vars.) is significantly represented, while aspen (*Populus tremuloides*), very abundant at present, is not represented in the sections because its pollen is poorly preserved.

Climatic trends are not clearly expressed, although in the Yukon Territory sections, the general increase of lodgepole from bottom to top suggests amelioration of climate in more recent time. Chronology is likewise indefinite. Radiocarbon dating of post-Wisconsin peat in the Anchorage area suggests an accumulation rate of 600 years per foot, which when applied to a section of this study near Anchorage, about 7 meters deep, indicates an age of more than 13,000 years. Another radiocarbon date for peat on the Kenai Peninsula indicates a depositional rate of about 800 years per foot which, when applied to the Anchorage section, denotes an age of over 18,000 years. The author believes that on the average the sections from interior Alaska and in the Yukon Territory represent an age up to 5000 years, while the deeper sections near the coast are somewhat older.

## INTRODUCTION

DURING the latter part of the summer of 1950, 75 peat sections were obtained from muskegs, bogs, swamps, and other sites of organic sedimentation in Alaska, the Yukon Territory, and British Columbia (fig. 1). The pollen profiles

* The author is grateful to the American Philosophical Society, the Geological Society of America, and the Arctic Institute of North America for generous grants to defray the expenses of the field work and equipment, and to the General Research Council of Oregon State College for grants for laboratory supplies and to support a research fellowship. Thanks are also due to Professor F. M. Beer, Oregon State College, and Professor R. F. Thaw, Corvallis High School, for assistance in collecting the peat

and other data derived from these sedimentary columns present a general picture of the forests that have existed in this vast region since the last glaciation, as well as some indication of climatic trends during the period represented. The pollen record as interred in these sections with that of 65 others along the Alaska Highway in British Columbia and farther south in Alberta provides a tentative history of the postglacial forests and climatic trends for western Canada and Alaska. Although this record is based upon only 140 sections taken along the principal highways, the relatively small number of forest tree species and the rather consistent and homogeneous structure and composition of the forests should permit a fairly representative picture of most of the region as indicated by the pollen profiles. Also the results of this study so far have presented to the author information and data which suggest how the problem should be continued on both more intensive and extensive scales.

Because this is a preliminary study and covers as much of the region in a general way as possible under the time limitation, the collection of sedimentary columns has been confined to sites largely adjacent to the main highways. Secondary highways are few and often impassable after rain, which occurs all too often during the late summer. As more side roads are constructed, the author hopes to continue with the problem.

## LOCATION AND CHARACTERISTICS OF THE SEDIMENTATION SITES

A transect of 36 sections was obtained along the Alaska Highway from the Yukon-British Columbia border to the Alaska-Yukon boundary, a distance of almost 600 miles (table 1). Five of these columns were taken in British Columbia between mileposts 733 and 786 where the highway swings south of the Yukon border. Two more sections were obtained in the Yukon on the Haines Highway, which connects Haines Junc-

sections, to Dr. Calvin J. Heusser, research fellow, for preparation of the peat for pollen analysis and most of the pollen identification, and to Dr. I. S. Allison, Chairman of the Geology Department, Oregon State College, for reading and criticizing the manuscript. The author is also grateful to the following members of the Alaska Terrain and Permafrost Section of the U. S. Geological Survey for reading the manuscript and offering valuable suggestions concerning permafrost, frost action and glaciation, as well as other phases of the paper: Troy L. Péwé, R. S. Sigafoos, R. F. Black, W. S. Benninghof, E. N. Muller, D. M. Hopkins, and E. T. Fernald.




Fig. 1. Generalized map of Alaska and the Yukon. Sites from which peat sections were obtained are shown by dots. Glacial boundaries from map by Capps, 1931. Modified from U. S. G. S. Alaska Map A.

tion at milepost 1016 on the Alaska highway with Haines, Alaska, at the north end of Lynn Canal, a distance of about 160 miles. Three other sections were procured along this highway in British Columbia.

Ten sections were obtained from muskegs along the Alaska Highway between the Yukon-Alaska border and Fairbanks, a distance of about 300 miles. Although the Alaska Highway technically goes only to Big Delta Junction at milepost 1420, it is now generally considered to extend all the way to Fairbanks. Actually, this latter segment is part of the original Richardson Highway that was constructed in 1920 from Fairbanks to Valdez, and it was the first highway from tidewater to the interior of Alaska. Two muskegs in the vicinity of Fairbanks also provide sections for this study.

Eight sites of organic sedimentation along the Richardson Highway were sampled between Delta Junction and its junction with the Glenn Highway at milepost 116 from Valdez, a distance of about 152 miles. Between Anchorage and the Richardson Highway, 9 muskegs and swamps were sampled along the Palmer and Glenn Highways, a distance of about 190 miles. Four sections were procured along the Slana-Tok Highway, which connects Tok Junction at milepost 1318 on the Alaska Highway with the Richardson Highway about 129 miles north of Valdez. The Slana-Tok Highway covers a distance of about 135 miles. The 75th column was taken from a muskeg in Alberta, about 75 miles north of Edmonton, and its pollen analytical data were included in a paper concerned with 10 other sedimentary columns in the Grande Prairie-Lesser Slave Lake region of Alberta (Hansen, 1952). The highways traveled and the approximate location of the sedimentation sites are shown on the map (fig. 1), and their locations are listed in table 1.

The pollen-bearing sections of this study were obtained from sites of several types of organic sediments. While the term "muskeg" has been applied to any swampy, marshy, boggy, or spongy area in the arctic and subarctic, as well as to tundra, there are several types of organic deposits formed under varying conditions and processes in these regions. The typical and perhaps the most widespread and extensive kind of deposit in this vast region, designated as muskeg, is that formed on flat and poorly drained terrain, and which has been built up

largely by one or more species of *Sphagnum* and ericaceous shrubs. Sedges also may be associated with the moss and shrubs and consist mostly of cottongrass (*Eriophorum* spp.) and *Carex*. Forbs and grasses also may be abundant although their presence and the species depend upon the amount of moisture present. This type of vegetative cover was apparently developed in the absence of ponded water and without the initial stages of hydrarch succession. This has been possible in a semi-arid region by the growth of *Sphagnum* on the surface in the presence of a high water table held up by permafrost. The accumulating vegetal material served as insulation which prevented the permafrost or seasonal frost table from receding downward in summer. This raised the water table higher, and with the absorbent plant remains acting as a wick, favorable conditions were maintained for continued organic deposition. The absence of pollen or aquatic plants in the lowest sediments as well as in the underlying clay and silt lends credence to this view (Hansen, 1949*a*, 1950*a*). The writer has suggested that the southern limit of muskegs formed by this process may approximate the southern limits of permafrost developed during and after the last glaciation (Hansen, 1952). In fact, it seems hardly possible that muskeg development could have been initiated in a region with an annual precipitation of less than 15 inches without the presence of permafrost except in lakes and ponds. Amelioration of the climate to a degree that would eliminate the permafrost also would possibly result in the extinction of many muskegs formed on flat terrain. Frozen substrata were noted farther south in muskeg than in inorganic substratum along the Alaska Highway, although this may merely emphasize the insulative qualities of the vegetative cover.

Shrubs commonly occurring on this type of muskeg include *Ledum groenlandicum*, *L. palustre decumbens*, *Andromeda polifolia*, *Chamaedaphne calyculata*, *Kalmia polifolia*, *Vaccinium oxycoccus*, *V. uliginosum*, *V. vitis-idaea minus*, *Empetrum nigrum*, *Betula glandulosa*, *Rubus chamaemorus*, and *Salix* spp. The most common tree is black spruce (*Picea mariana*). Tamarack (*Larix laricina*) was noted on bogs and muskegs as far as milepost 688 in the Yukon, and again in the Fairbanks region. An occasional lodgepole pine (*Pinus contorta latifolia*) was noted on muskegs in the Yukon, with an abundance observed on a muskeg at milepost 666. On some of the

TABLE 1

Peat Sections from the Yukon Territory

| Miles from Dawson Creek, B. C. | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| 634—Watson Lake | | | |
| 647 | 5.7 | 5.6 | Lake; marly; pine, spruce, birch, aspen adjacent |
| 655 | 1.0 | 1.0 | Sedge, ericads; pine, spruce adjacent |
| 664 | 2.8 | 2.5 | Sedge; pine, tamarack, spruce adjacent |
| 666 | 4.5 | 4.2 | Muskeg; spruce, pine, tamarack, ericads |
| 681 | 1.5 | 1.5 | Sedge bog; spruce, tamarack on bog |
| 688 | 1.5 | 1.5 | Muskeg; spruce, tamarack, pine on bog |
| 699 | 1.5 | 1.5 | Sedge, ericads, juniper; pine, spruce adjacent |
| 707 | 1.9 | 1.5 | Muskeg; spruce, ericads; pine adjacent |
| 716 | 2.0 | 2.0 | Lake; sedge, ericads; pine, spruce adjacent |
| 724 | 2.0 | 2.0 | Muskeg; spruce, ericads |
| 737—British Columbia | 2.9 | 2.9 | Muskeg; spruce, ericads |
| 748 | 0.7 | 0.7 | Muskeg; burned |
| 759 | 1.8 | 1.3 | Muskeg; spruce, ericads |
| 770 | 3.5 | 3.5 | Muskeg; pine, spruce adjacent |
| 779 | 2.0 | 2.0 | Lake; pine, spruce adjacent |
| 796—Yukon | 1.0 | 0.8 | Lake; sedge; pine, spruce adjacent |
| 840 | 1.7 | 1.6 | Sedge-sphagnum bog; pine adjacent |
| 859 | 4.9 | 3.6 | Lake; sedge, sphagnum |
| 872 | 5.5 | 4.4 | Sedge-sphagnum bog; pine, spruce adjacent |
| 899 | 1.0 | 0.9 | Pond; sedge; pine, spruce |
| 918—Whitehorse | | | |
| 923 | 2.0 | 0.6 | Sedge-sphagnum bog |
| 931 | 0.8 | 0.4 | Muskeg; ericads; spruce, pine adjacent |
| 947 | 1.4 | 0.1 | Pond; sedge; sandy area |

TABLE 1 (cont.)

Peat Sections from the Yukon Territory

| Miles from Dawson Creek, B. C. | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| 974—Champagne | | | |
| 986 | 0.5 | 0.1 | Sedge bog; spruce, aspen adjacent |
| 1009 | 1.0 | 0.4 | Sedge-hypnum bog; spruce, willow adjacent |
| 1013 | 1.5 | 1.1 | Pond; sedge; spruce, willow adjacent |
| 1016—Haines Jct. | | | |
| 1037 | 0.7* | 0.7 | Lake; sedge; spruce adjacent |
| 1045 | 0.7 | 0.7 | Muskeg; spruce, ericads |
| 1102 | 0.6 | 0.4 | Muskeg; spruce, ericads |
| 1128 | 0.7 | 0.6 | Sedge; burned |
| 1164 | 1.0 | 0.8 | Sedge; spruce, ericads |
| 1170 | 0.52* | 0.5 | Muskeg; spruce, ericads |
| 1189 | 0.5 | 0.4 | Lake; spruce, ericads |
| 1192 | 2.0 | 1.8 | Lake; sedge, spruce, ericads |
| 1200 | 0.52* | 0.4 | Muskeg; spruce, ericads |
| 1221 | 0.52* | 0.5 | Muskeg; spruce, ericads |

*Permafrost

TABLE 2

Peat Sections between Haines Junction, Yukon, and Haines, Alaska

| Miles from Haines, Alaska | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| 72 B. C. | 1.0 | 0.9 | Sphagnum; spruce, juniper |
| 79 B. C. | 0.7* | 0.6 | Sedge-sphagnum; willow; above timberline |
| 84 B. C. | 1.0 | 0.7 | Sedge-sphagnum; few spruce on hillsides |
| 103 Yukon | 0.52* | 0.5 | Sedge-sphagnum; willow, spruce scattered |
| 135 Yukon | 0.5 | 0.1 | Sedge-sphagnum; spruce adjacent |

TABLE 3

Peat Sections along the Alaska Highway from Yukon Border to Fairbanks

| Miles from Dawson Creek, B. C. | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| 1248 | 0.52* | 0.4 | Muskeg; ericads, spruce |
| 1270 | 0.5* | 0.5 | Muskeg; ericads, spruce |
| 1297 | 0.52* | 0.52 | Muskeg; ericads, spruce |
| 1299 | 0.52* | 0.52 | Muskeg; ericads, spruce |
| 1318—Tok Junction | | | |
| 1336 | 0.5* | 0.5 | Muskeg; spruce |
| 1353 | 0.5* | 0.5 | Muskeg; spruce, ericads |
| 1387 | 0.9* | 0.9 | Muskeg; spruce, ericads |
| 1390 | 0.6* | 0.6 | Muskegs; ericads, spruce |
| 1420—Big Delta Junction | | | |
| 1463 | 0.5* | 0.5 | Muskeg; ericads, tamarack, spruce |
| 1488 | 0.6* | 0.5 | Muskeg; ericads, tamarack, spruce |
| 1523—Fairbanks (near) | 2.0* | 1.9 | Muskeg; ericads, spruce |
| Fairbanks (near) | 2.3 | 1.6 | Small pond; aquatics, sedge, willow |

*Permafrost at lowest level

drier and better-drained muskegs fire may favor an invasion of paper birch (*Betula papyrifera* var.), reindeer moss (*Cladonia* sp.), and hairy cap moss (*Polytrichum* sp.). Bunchberry (*Cornus canadensis*) also seems to be more abundant on burned muskegs. This type of muskeg becomes more shallow northward, and in the Tanana River valley; most of them sampled are about 0.5 meter deep (table 1). Farther south in British Columbia muskegs formed without evidence of an initial aquatic stage may be more than 5.0 meters deep (Hansen, 1949*a*, 1950*a*, 1952). Permafrost or a seasonal frost residuum was encoun-

tered in all muskegs sampled between the Yukon-Alaska border and Fairbanks, and southward in the Yukon to milepost 1037. Frost also was noted in muskegs along the Alaska Highway in British Columbia (Hansen, 1950*a*). In muskegs with a frozen substratum it was found that a thicker section of organic sediments could be obtained between the mounds than through them. The larger mounds, in which ice was encountered a foot or so from the surface, may be similar in structure and development to the "palsas" in Labrador described by Wenner (1947), in which almost pure ice occurs in large mounds covered with an insulating layer of *Sphagnum*.

Another type of site from which sections were obtained is the "niggerhead" muskeg which is characterized by tussocks or hummocks of one or more species of cottongrass (*Eriophorum* spp.). The tussocks may be as high as 18 inches from the crown of the roots down to the substratum between the columns of cottongrass roots. Standing water may be present between the hummocks or other plants may grow on the damp substratum between. *Hypnum* spp. frequently occurs between the hummocks, and bog birch and blueberry are also abundant on the "niggerhead" muskeg. The sediments in this type of muskeg were found to be shallow and silty, with little organic matter. Pollen analysis also reveals that little pollen is preserved in sediments accumulated under this type of vegetative cover.

Many of the sedimentary columns were obtained from sites of hydrarch succession that were initiated in lakes, ponds, sloughs, or in valleys with poor drainage. In most cases open water is still present with well-defined concentric zones of vegetation representing various stages of succession. Some support typical bog seres and culminate in Sphagnum-Ericad-black spruce cover. Many of them, however, apparently are progressing toward a sedge-willow-bog birch cover, especially in the Takhini Valley beyond Whitehorse and southeast of Whitehorse in the drier, parkland areas. The lower sediments in these muskegs often consist of marly deposits indicating an abundant source of calcium carbonate in the early aquatic stages of succession. Aquatic plants growing in some of these ponds are encrusted with calcium carbonate. The thickest organic sections were procured from bogs that have developed in ponded water, although in the lower levels pollen is often scarce. Permafrost was not found in sediments that accumulated in lakes, even in areas where it was encountered in the typical muskeg.

TABLE 4

Peat Sections along the Richardson Highway from Big Delta Junction to Glenn Highway

| Miles from Fairbanks | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| 119 | 0.6 | 0.5 | Small pond; willow, birch, alder adjacent |
| 126 | 0.5 | 0.5 | Moss, sedge, on slope |
| 167 | 1.0 | 0.5 | Sedge, willow, sphagnum, bog birch |
| 192 | 1.0 | 0.5 | Small lake; spruce, ericads |
| 196 | 1.5 | 1.3 | Small lake; sedge |
| 210 | 1.1* | 0.7 | Muskeg; ericads, bog birch, spruce |
| 225 | 1.0 | 0.8 | Lake; muskeg, ericads, spruce, bog birch |

TABLE 5

Peat Sections between Anchorage and Junction of Glenn and Richardson Highways

| Miles from Anchorage | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| Near city airport | 0.6 | 0.5 | Muskeg; spruce, bog birch, ericads, sundew, sphagnum |
| 15 | 6.9 | 5.2 | Lake; floating stage, ericads, white birch, bog birch |
| 26 | 0.4 | 0.4 | Muskeg; spruce, ericads |
| 28 | 4.5 | 2.6 | Wet; spruce, ericads, bog birch |
| 88 | 3.0 | 2.7 | Wet; spruce, ericads, bog birch, buckbean |
| 122 | 0.6 | 0.5 | Muskeg; spruce, ericads |
| 141 | 0.5 | 0.5 | Lake; ericads, sedges |
| 151 | 2.0 | 1.8 | Lake; ericads, sedges, floating stage |
| 158 | 3.9 | 3.0 | Muskeg; ericads, sedge, bog birch |

TABLE 6

Peat Sections along Slana-Tok Highway between Richardson Highway and Tok Junction

| Miles from Tok Junction | Total depth of section in m. | Depth of pollen profiles in m. | Characteristics of site |
|---|---|---|---|
| 105 | 0.9 | 0.3 | Lake; ericads, bog birch, willow |
| 95 | 0.55 | 0.55 | Muskeg; spruce, ericads; dry |
| 65 | 3.0 | 1.4 | Pond; limnic peat; Hypnum |
| 36 | 2.0 | 0.9 | Pond; floating stage; willow, alder, bog birch |

The three types of sedimentary sites described above represent in general the source of most of the pollen-bearing sediments of this study. Each site differs in physical and floristic detail according to the immediate physiographic and edaphic conditions and the drainage pattern. Inasmuch as the sediments are merely a means to an end, namely, the pollen record interred therein, further detail here is of little significance to the problem at hand.

The thickest section, 6.9 meters, was obtained from a site adjacent to a lake about 15 miles inland from Anchorage. The typology reveals a typical hydrarch succession with the lowest horizons consisting of fine, colloidal, limnic sediments, grading upward into fibrous peat. The submerged and floating stages in the lake are somewhat similar to those in the Puget Lowland of western Washington (Hansen, 1947). No permafrost was encountered, and the section extends down to coarse, impenetrable gravel. In those muskegs where a frozen substratum was present, in most instances the frost table had receded downward during the summer to the inorganic sediments underlying the fibrous peat. In obtaining sections, the permafrost was penetrated at least a decimeter or more. Little or no pollen was noted in those sediments that had been frozen, except in places where they included the lower part of the organic material.

Five muskegs between milepost 1128 in the Yukon and 1270 in Alaska were underlain with pumice, but the source is unknown. Strata of volcanic glass, indiscernible to the naked eye

but evident under the microscope, also occur in many of the sections along the Alaska Highway. Two of these ash horizons seem to be chronologically equivalent in the several sections in which they occur (figs. 2-7). One of these volcanic ash strata may have had its source in the St. Elias Range to the southwest. Capps (1916) assigned glass in the Fairbanks region to this source and set the date of the eruption about 1400 years ago. Volcanic glass has been noted by Heusser (1952) in bogs along the southeastern coast of Alaska. In the Sitka region, glass has been assigned to an eruption of Mt. Edgecombe 20 miles distant, and upon the basis of its stratigraphic position and the recorded forest succession has been dated at about 5500 years. Ash in the upper six inches of peat sections along the Alaska Highway in northeastern British Columbia has been ascribed to the eruption of Mt. Katmai in 1912, although Heusser finds no evidence of ash attributable to this source in the muskegs of southeastern Alaska. Kerr (1936) notes that postglacial volcanic activity occurred in mountains 50-60 miles east of Wrangell, possibly 100 years ago. Undoubtedly postglacial volcanic activity has occurred many times in the Coast Range of northern British Columbia and the mountains of Alaska, and it does not seem possible to assign the volcanic glass and pumice that occur in the sedimentary columns to any particular source. They may, however, serve as valuable chronological markers in correlating the recorded forest succession from muskeg to muskeg and from one area to another, when they can be identified.

The Alaska Highway has its southern terminus at Dawson Creek in eastern British Columbia about 12 miles from the Alberta-British Columbia border. It extends north and somewhat west through the rolling plains and the eastern Rocky Mountain foothills to Muskwa near Fort Nelson. It then turns almost due west and cuts through the Rocky Mountains and onto the Liard Plain (Bostock, 1948). The highway then follows up the Liard valley nearly to Watson Lake in the Yukon and continues westward across the Dease Plateau and the northern part of the Cassiar Mountains. On the Teslin Plateau it follows lake and river valleys to Whitehorse, the Takhini Valley and Shakwak Valley northwestward almost to the Yukon-Alaska border.

In the Yukon the Alaska Highway lies within the limits of Wisconsin glaciation, although it has not been determined to

what extent the Late-Wisconsin ice covered this region (Flint, 1945, 1947; Denny, 1952). The last 25 miles before the Yukon-Alaska boundary, however, apparently were not covered by Wisconsin ice. Also less than 50 miles to the north of the highway in the Yukon River drainage there is a vast unglaciated region extending nearly 200 miles eastward into the Yukon and reaching northward almost to the Arctic Ocean (fig. 1).

The highway a short distance beyond the Yukon-Alaska border enters the Tanana River valley which it follows to Fairbanks. To the south of the Tanana River is the Alaska Range and to the north lies the interior plateau extending northward to the Brooks Range and drained by the Tanana and Yukon rivers. This area was not glaciated by Wisconsin ice because of low precipitation, and the poorly nurtured glaciers on the north slope of the Alaska Range were unable to flow out upon the plateau (Capps, 1931). Although the muskegs along the highway between the border and Fairbanks lie in an unglaciated region, their sites were undoubtedly modified by glacial meltwater from the Alaska Range which was carried westward by the Tanana River drainage.

The Richardson Highway, south of Big Delta, cuts through the Alaska Range following along in valleys and on tablelands. The Glenn Highway follows westward from its junction with the Richardson between the Talkeetna Mountains to the north and the Chugach Mountains to the south. It then extends down the Matanuska Valley to Palmer, then the Knik River to tidewater and along the Knik Arm of Cook Inlet to Anchorage. The Slana-Tok Highway follows through rugged country between the Wrangell and Mentasta Mountains and then emerges upon a flat terrace in the Tanana River valley for the last 25 miles to Tok Junction on the Alaska Highway. The muskegs and bogs along these highways that provide sedimentary columns for this study lie on sites that were glaciated by the Late Wisconsin ice and modified by meltwater during its recession.

### FORESTS ALONG THE ALASKA HIGHWAY IN THE YUKON

The forests along the Alaska Highway in the Yukon Territory are comprised of nine arboreal species. These are white spruce (*Picea glauca*), black spruce (*P. mariana*), alpine fir (*Abies lasiocarpa*) lodgepole pine (*Pinus contorta latifolia*),

tamarack (*Larix laricina*), aspen (*Populus tremuloides*), balsam poplar (*P. tacamahacca*), black cottonwood (*P. trichocarpa*), and paper birch (*Betula papyrifera* vars.). Sitka spruce (*Picea sitchensis*) occurs sparingly in southwestern Yukon near the Haines Highway. Phytosociologically only two of these, white spruce and alpine fir, are considered to be permanent or climax species. The others owe their persistence to either immature succession or disturbance such as fire, cutting, landslides, new floodplain deposits, rabbit depredations, and other disrupting factors. Fire, of course, has been by far the most significant and important factor in the persistence of the nonclimax species. It is evident that most of the area along the highway has burned recently or not far in the past. There is every stage of succession in evidence, resulting from the complete destruction of the forest over many square miles to partial destruction and small burns. Lodgepole and aspen have benefited most from fire, while white birch and the poplars may be favored under certain conditions. Tamarack is confined to muskegs and is only locally abundant. Black spruce is also chiefly a muskeg dweller, but the vast extent of the muskeg in this region and its lower susceptibility to fire than the upland forest permit this species to persist indefinitely in the absence of competition. Under the present climate and drainage it seems that the muskegs are permanent and black spruce may be considered as a physiographic climax. White spruce prefers the better-drained slopes and uplands, although it also thrives on floodplains where the soil is light. The poplars also occur on floodplains and sand bars where they may form almost pure stands. An understory of white spruce in the older stands, however, indicates the eventual replacement of the poplars. Paper birch is not abundant and rarely forms anything resembling pure stands. It often invades the drier muskegs after they have been burned, but it is most common on well-drained slopes where fire has occurred in the past; its relation to other species is not clear. Alpine fir is not common adjacent to the highway, but it occurs on the upper slopes and becomes the dominant tree at timberline at about 4000 feet (Halliday, 1937).

Of the nine arboreal species in the Yukon, lodgepole pine and alpine fir are Cordilleran in range. Lodgepole has a vast range, extending from Lower California to central Yukon and from the Pacific Coast eastward to the east slope of the Rocky

Mountains. In Canada it ranges eastward in Alberta to Smith and Edmonton and it also occurs as an outlier in the Cariboo Mountains south of Great Slave Lake (Raup, 1935). Alpine fir also extends eastward to central Alberta (Halliday and Brown, 1943). White and black spruce, balsam poplar, and tamarack are species of the boreal forest that extends in a broad band from the Gulf of St. Lawrence northwestward across Canada to Alaska, with the first three extending southward in the Rocky Mountains. Two other species of the boreal forest, jack pine (*P. banksiana*) and balsam fir (*A. balsamea*) range westward to north central Alberta, the former overlapping the range of lodgepole pine for 200 miles or more (Moss, 1949). Paper birch is also more or less coextensive with the spruces, but it breaks up into several varieties in the northwestern territories and Alaska. It is reasonable to assume that any species with as great a range as lodgepole pine must consist of a large number of biotypes, which is responsible for its wide ecologic amplitude.

The first species observed to drop out northward along the Alaska Highway is tamarack which was last noted in a muskeg at milepost 688, although it may extend farther at points remote from the highway. The only other arboreal species to reach the end of its range along the highway is lodgepole, which was last noted about 25 miles beyond Whitehorse at milepost 945. It is reported to occur as far northwest as Kluane Lake. The only portion of the Alaska Highway between Watson Lake and the Alaska-Yukon border that does not normally support forests is in the Takhini-Dezadeash valleys west of Whitehorse for a distance of about 100 miles. Here the soil is light and in some places sandy, with dune areas in the vicinity of Champagne. The vegetation has a natural parkland appearance, with scattered stands of white spruce, aspen, willow, and open grassy areas with an abundance of forbs. In the Kluane Lake region and westward almost to the Alaska border the forest consists almost entirely of white spruce, although black spruce was again noted in the White River valley and is also common in the Tanana River valley in Alaska.

In general, it can be said that the principal forest mixture or type along the Alaska Highway in the Yukon is white spruce, lodgepole pine, and aspen. Between Watson Lake and Whitehorse, pine forms vast pure stands on the lighter soils and in

other areas where fire has been frequent. White spruce usually predominates on the heavier clayey soils in well-drained upland areas, while black spruce usually occupies muskeg on the valley floors, although it may also occur in upland communities of white spruce and pine, especially on northern exposures. It appears that aspen is a fire species that prospers after several frequently recurring fires that first kill the spruce and/or pine and then their reproduction. Aspen, growing faster and sprouting by means of root suckers, thrives until it is replaced first by pine and then by spruce if the fireless interim is sufficiently long.

## FORESTS OF ALASKA

Alaska has an area of 586,400 square miles, and its vegetation may be broadly classified as coastal forests, interior forests, and treeless tundra and grassland. The coastal forests of southeastern Alaska cover about 10 per cent, the lightly timbered interior forests about 60 per cent, and the nontimbered tundra perhaps 30 per cent (Heintzelman, 1951). These figures are merely estimates and differ from those of other observers of the region. Within each of these formations there are many localized types because of physiography, soil, climate, and fire.

The coastal forests of southeastern Alaska, from which there are no peat sections for this study, represent an extension of the coastal forests of Oregon, Washington, and British Columbia. These are comprised of Sitka spruce (*Picea sitchensis*), western hemlock (*Tsuga heterophylla*), western red cedar (*Thuja plicata*), lodgepole pine, and silver fir (*Abies amabilis*). In addition, at higher elevations and under favorable conditions at sea level such species as Alaska cedar (*Chamaecyparis nootkatensis*), mountain hemlock (*T. mertensiana*), and alpine fir are present. The northernmost extension of the Coast Forest reaches Cook Inlet, with Sitka spruce ranging slightly farther north than western hemlock and Alaska cedar. Silver fir (*A. amabilis*) occurs at the southern end of the Alaskan coastal forest (Taylor, 1950; Munns, 1938). Broadleaf arboreal species in southeastern Alaska are paper birch (*Betula papyrifera* vars.), red alder (*Alnus rubra*), thinleaf alder (*A. tenuifolia*), Stika alder (*A. sinuata*), Douglas maple (*Acer Douglasii*), and northern black cottonwood (*Populus trichocarpa*). Several species of willows of tree stature are also present.

Since 1750 the valley glaciers in southeastern Alaska have been receding (Lawrence, 1950), affording deglaciated terrain open for invasion by the vegetation of the region. The general trend of succession seems to be initial invasion by alder and willow, and cottonwood, followed by Sitka spruce which largely replaces the less tolerant broadleaf species, with finally a climax of western hemlock and Sitka spruce. Lodgepole seems to be restricted largely to muskegs, while Alaska cedar also occurs in such habitats in addition to its occurrence at timberline with alpine fir and mountain hemlock. Pollen analysis of peat sections from southeastern Alaska has been carried out by Heusser (1952), and the results pertinent to this study will be discussed later.

The interior forests of Alaska are composed of a limited number of arboreal species. These are white and black spruce, paper birch, aspen, balsam poplar, tamarack, and alpine fir. Although maps by Harlow and Harrar (1941) and Munns (1938) show the range of lodgepole as extending into eastern Alaska from the Yukon, Taylor (1950) states that this species does not enter Alaska from the Yukon, and extends northward along the coast to the head of Lynn Canal at Skagway and Glacier Bay. No lodgepole was noted in interior Alaska along the highways, although this, of course, does not preclude its existence in remote areas. Alpine fir likewise is of rare occurrence in interior Alaska but it was noted occasionally along the Slana-Tok Highway between Sinona and Mentasta at higher elevations.

It is generally conceded by botanists, and others who have made observations of the forests in this region, that white spruce is to be considered as the climatic climax dominant, and that in the absence of fire, disease, animal depredations, and other disturbances for several centuries, the upland stands of timber would consist largely of white spruce. This, unfortunately, has not been the case, and it is estimated that during the past half century the average area burned annually has been at least 1,000,000 acres and so not more than 20 per cent of the original white spruce forest remains (Lutz, 1951). The criteria for considering white spruce as the climax species or as the culmination of successional development on the uplands are the facts that it attains greater age than the other boreal species with which it is associated, it is more tolerant of shade, it is less

subject to disease and decay at an early age, it more readily germinates and develops on organic substratum, and it grows more rapidly than its associates on the well-drained, upland soils. Severe fires in white spruce stands usually result in its total destruction and succession begins anew with subclimax species. The composition of the new young forest depends upon the species that survive in proximity to the burn. Birch, aspen, and balsam poplar are the usual invaders, although the relative success of these species depends upon the physiographic and edaphic factors. If the organic material in the soil is destroyed, aspen will be favored; whereas if the litter and duff have not been destroyed spruce and birch, if seed is available, will invade and successfully compete with aspen during this early stage. Undisturbed succession, however, will culminate in white spruce and birch, and if continued for a century or more pure spruce will obtain. The muskegs and other poorly drained areas support essentially pure stands of black spruce. Tamarack occurs with black spruce in the valleys of the Tanana, lower Yukon, upper Kuskokwim, and upper Koyukuk rivers. Along the Alaska Highway it first was noted near Big Delta Junction. While black spruce is a seral species of the latter stages of hydrarch succession on muskegs and bogs farther south, it may be considered as a physiographic climax in interior Alaska, because of its evident permanence. Black spruce readily reproduces vegetatively by layering. The permanency of black spruce is further assured by the absence of fire on the wet muskegs, whence seed is provided for the adjacent burned areas. Thus black spruce has been gradually invading uplands following fire in white spruce. Once established in these areas it has the advantage of serotinous cones, which remain closed and are retained for several years on the trees. Following fire the cones are opened and the seeds dispersed in great quantities so that a black spruce stand may often regenerate to black spruce. This process is similar to that of lodgepole in the Rocky Mountains and Douglas fir in parts of the Pacific Northwest which are maintained through frequently recurring fires. Repeated severe burning may cause reversion from black spruce to earlier stages consisting of alder, dwarf birch, and willow.

One of the important basic factors in forest distribution and composition in interior Alaska as well as in the Yukon is permafrost. The interaction between permafrost, slope, expo-

sure, soil texture and fire plays a critical role in determining the vegetative cover. Destruction of the litter and duff may also lower the permafrost table, permitting trees with deeper root systems to thrive. Elimination of the organic material in the soil and lowering of the permafrost table may also lower the water table and the drier conditions may encourage less mesophytic species. Aspen is particularly favored under these conditions because it prefers a mineral soil for germination, deep and well-drained light soils, and south slopes where it simulates the climax situation. Absence of white spruce in the understory may indicate that the permafrost table is several feet below the surface and the soil too dry (Stoeckler, 1949). The ability of aspen to sprout from roots after a fire also favors its persistence even under less favorable conditions.

Balsam poplar prefers deep, moist, light soils and occurs chiefly on newly formed floodplains and river bars, although it also invades the uplands where mineral soil is exposed. It has a deeper root system than white spruce and its presence indicates a permafrost table several feet below the surface. Birch germinates best on mineral soil, but it can grow under a wide range of moisture conditions. It has a shallow root system but prefers deeper soil than does white spruce. Although it invades burned areas with spruce, it is much less tolerant and in time is replaced by spruce.

The mosaic of forest types in the interior of Alaska is not well understood, but it seems obvious that the thermal regime of the soil as influenced by slope, exposure, drainage, depth and structure of soil, and fire in relation to the ecological requirements of the trees is one of the more significant controlling factors.

## POSTGLACIAL FORESTS

In considering the postglacial forests as portrayed by the pollen record for the time represented by the pollen-bearing columns, the usual sources of error inherent in pollen analysis should be kept in mind. Perhaps the most significant in the region of this study is the evident failure of aspen and cottonwood pollen grains to be preserved in the sediments. These species, especially the former, have undoubtedly been important in forest succession, particularly in relation to fire. Although it seems evident that fire has been far more extensive and destructive during the past half century than perhaps during the




**Fig. 2. Pollen diagrams from along the Alaska Highway in the Yukon Territory from milepost 647 to 666.**




Fig. 3. Pollen diagrams from along the Alaska Highway in the Yukon Territory from milepost 681 to 707.

rest of postglacial or even Pleistocene time, fires started by early man and lightning probably have favored the persistence of aspen in abundance for many millennia. The absence of aspen pollen in the sediments, therefore, causes the picture of postglacial forests to be incomplete and also results in the overrepresentation of those species that are recorded. Thus most of the pollen profiles from Alaska, especially in the interior, reveal nearly 100 per cent of spruce. Although these pollen profiles may seem to be of no significance, they reveal whether or not certain species such as lodgepole pine and alpine fir ever have existed in interior Alaska during the Postglacial.

Lodgepole is recorded as having been important in postglacial forest succession northwestward to milepost 931 in the




Fig. 4. Pollen diagrams from along the Alaska Highway in the Yukon Territory from milepost 716 to 759.

Yukon. This section is located about 15 miles within the northwestern limits of lodgepole as noted along the highway. Lodgepole is also appreciably represented in sections along the Haines Highway, the pollen of which undoubtedly was carried eastward from the coastal area of southeastern Alaska. The record of lodgepole does not become progressively weaker northwestward toward the end of its range as one might expect. Rather it is strongly represented in the section at milepost 931, beyond which it abruptly ceases to be recorded appreciably in any part of the column (figs. 2-6). In the first 22 sections along the highway there are two general patterns of lodgepole-spruce relationships exhibited. The more common is a general increase of lodgepole from low proportions at the bottom to 60 per cent or more, and in some cases to 80 per cent, in the middle third of the profile and maintenance of this preponderance to the top (figs. 2, 4, 5). In two instances lodgepole supersedes spruce at the bottom and then declines before it attains its strong predominance in the upper half or two-thirds (fig. 2). This deviation may represent an earlier stage not recorded in the other sections as it is expressed in only two of the deeper sections, 5.6 and 4.0 meters at mileposts 647 and 666 respectively. Another section, 4.0 meters, at milepost 872 exhibits an increase of pine to 90 per cent or more in the upper third (fig. 6). The other pattern reveals lodgepole as strongly predominant throughout the column, with proportions of mostly 75 per cent or more from bottom to top. The profiles of spruce are almost the exact converse of those of pine, except in those sections where grass is appreciably represented. Alpine fir is represented consistently but in low proportions throughout most of the sections. The maximum of this species is only 10 per cent at milepost 872 (fig. 6). Spruce pollen profiles are based upon both black and white spruce, which reflect the influence of the forest on the muskeg and adjacent poorly drained areas. This is partially offset by the occurrence of black spruce in uplands where repeated conflagration has destroyed the white spruce forests so that seed source of this species is too far removed for its immediate reinvasion. The occurrence of black spruce on the muskeg may be compensated for by a limited amount of pine also thereon, and the latter's greater pollen production. This is evinced by the overwhelming preponderance of lodgepole pine in most of the sections, espe-




**Fig. 5. Pollen diagrams from along the Alaska Highway in the Yukon Territory from milepost 770 to 858.**




Fig. 6. Pollen diagrams from along the Alaska Highway in the Yukon Territory from milepost 872 to 1037.

cially in the upper half. The average number of horizons of pine predominance in the sections to milepost 931 is much greater than that of spruce, and while the coarser and more fibrous nature of the peat in the upper levels probably represents more rapid deposition, the average portion of the profiles denoting pine predominance suggests that it constitutes more than half of the time represented. The occurrence of two strata of volcanic glass within the period of pine predominance in most sections suggests synchronism of the pine record.

Grass is well represented in the first four sections northwestward as far as milepost 666, and then it becomes sporadically recorded in the next five sections to milepost 724 (figs. 2, 3, 4). It is poorly but almost continuously represented in the next four sections to milepost 770, and then becomes absent except for an occasional pollen grain in the other sections of this study. The grass record is somewhat anomalous in that it is not represented in the sections obtained in the parkland areas west of Whitehorse where grasses are significantly abundant. While grass is strongly represented in the first four sections and indicates some environmental influence, the lack of a regional consistency does not permit a tenable interpretation. Ordinarily grass proportions as high as 15 per cent suggest drying and/or warming in a forested region. The




Fig. 7. Pollen diagrams from along the Alaska Highway in the Yukon Territory from milepost 1045 to 1192.

muskegs from which the sections were obtained did not support grasses and most of them surrounded open water. The adjacent areas likewise do not support grasslands. Uplands remote from the site of the sediments were not observed, however, and grass pollen may have come from such areas. There does not seem to be any correlation of the grass profiles with those of pine and spruce, especially in relation to fire in recent time which has apparently favored pine and aspen. In fact, most profiles reveal absence or decline of grass in the upper levels.

Sedimentary columns obtained from sites beyond milepost 931 show an abrupt drop in the pine proportions to only a few per cent (figs. 6, 7). Pine is well represented in five sections taken between Haines Junction at milepost 1016 and Haines, Alaska. Three of the thicker sections reveal pine proportions ranging from 1 to 42 per cent, with an average of over 20 per cent (fig. 8). The forests at this elevation are sparse and no pine was noted, so its pollen must have been carried up and inland from the coastal forests. Western hemlock is also appreciably represented in these columns, which with the presence of some mountain hemlock pollen further indicates the effect of winds moving inland from the coast during the period of anthesis. Alpine fir, which becomes absent in the profiles at milepost 1037, is sparingly represented in some of these sections. Some of the spruce pollen is undoubtedly Sitka spruce from the coast and the few scattered trees along the highway.

Pollen profiles from 17 sedimentary columns along the Alaska Highway in northeastern British Columbia from Dawson Creek




Fig. 8. Pollen diagrams from along the Haines Highway, British Columbia. Mileposts are from Haines, Alaska.

to Watson Lake reveal trends of forest succession both similar to and different from those of this study (Hansen, 1950*a*). In six of these sections spruce is predominant throughout, and pine in only three. Similarity of trends is expressed by pine expansion in the upper levels to supersede spruce at the top in 10 columns. In general spruce has apparently been more abundant along the first 600 miles of the Alaska Highway in British Columbia than in the Yukon to milepost 1009. In a 6.6-meter section at milepost 253, spruce is recorded to 80 per cent or more throughout its entire profile. Still farther south and east in the Grande Prairie-Lesser Slave Lake region, spruce and pine are more nearly equally represented in most of 11 sections, although pine increases in the upper horizons to supersede spruce in all of them (Hansen, 1952). Still farther south in the vicinity of Edmonton and near the southern limits of spruce and pine in central Alberta, pollen profiles from seven muskegs show pine predominance at the bottom in six of the sections and throughout generally (Hansen, 1949*a*). An increase in pine at the top in six of the profiles is consistent with its record in most sections to milepost 931, suggesting the influence of fire since the advent of white man, perhaps further favored by a drier climate during the past 200 years. West of Edmonton and nearer the Rocky Mountains a similar pattern for pine is shown in three out of four profiles (Hansen, 1949*b*). In general, spruce becomes more consistently predominant in the lowest pollen-bearing level progressively northwestward and persists in preponderance longer, but in more recent time lodgepole pine has expanded over the region represented in the pollen profiles because of more favorable conditions in the past few centuries. Why its record should so abruptly become absent beyond milepost 931 is difficult to explain unless this transition zone also constitutes the southern limits of permafrost which is slowly moving northward in response to a warming climate.

The forests as recorded in sections taken between the Yukon border and Fairbanks become more and more preponderantly spruce, so there is no need for profiles to be illustrated. A few pine pollen grains occur sporadically in sections as far as milepost 1390, about 135 miles south of Fairbanks. Two sections near Fairbanks and the first two south of this city contain a considerable amount of alder and birch pollen in the upper horizons while birch is also well represented near the bottom.

Much of the birch pollen is probably from paper birch on uplands while some is from bog birch on the muskeg. Dwarf birch may also be represented as this species is common on oft-burned areas where the arboreal cover has been entirely depleted. Volcanic glass occurs in the upper two decimeters of sediments, and another layer is also present near the bottom in most of these sections.

In the four sections procured along the Slana-Tok Highway southwest of Tok Junction, spruce is recorded to practically 100 per cent throughout the pollen-bearing strata. Only two pollen grains of pine and one of fir were noted. The thickest section, 3.0 meters, which was taken from the border of a pond in a deep depression, reveals pollen only to 1.6 meters, while the sediments down to 2.5 meters consist of colloidal limnic peat of largely organic matter. The absence of tree pollen suggests that forests did not exist in the region when these sediments were laid down. The adjacent hillsides support some birch, and yet very little birch pollen was noted. In all sections volcanic glass occurs in the upper two decimeters and again near the bottom, as in sections along the Alaska Highway between the Yukon border and Fairbanks.

Sedimentary columns taken from eight sites along the Richardson Highway south of Big Delta Junction to the Glenn Highway reveal that spruce is recorded to 100 per cent throughout their pollen-bearing strata. A single pine pollen grain was observed in the uppermost level of a section taken about 19 miles south of Delta Junction. Two sections taken in this area show an abundance of alder and birch pollen throughout and some willow and sedge pollen at a few of the horizons. That much of the adjacent area is covered with alder and dwarf birch explains the source of the pollen. A section taken from a willow-sedge muskeg on Isabella Pass, at an elevation of about 3200 feet, has little pollen throughout the silty sediments but spruce is recorded to 100 per cent at all levels. In the other sections, the last of which was taken about 122 miles south of Big Delta Junction, the few pollen grains of mountain hemlock noted probably were carried inland from the coast. In most columns along the Richardson Highway the lower sediments are composed largely of silt, and there is little or no pollen present in the lowest 2-4 decimeters so as to suggest the absence of forests in adjacent areas when they were deposited.

Nine sections along the Palmer-Glenn Highway, a distance of about 190 miles, reveal a preponderance of spruce in those from higher elevations inland, while in those situated in the lowlands within 30 miles of Anchorage birch is strongly represented (fig. 9). The presence of paper birch on the adjacent slopes accounts for the abundance of its pollen in the upper levels, and it can be presumed that birch has been an important component of these forests during most of the time represented by the sediments. In the thickest section of this study, 6.9 meters, about 15 miles east of Anchorage, birch shows a stronger




Fig. 9. Pollen diagrams from along the Palmer-Glenn Highway, Alaska. Mileposts are from Anchorage.

average representation than does spruce. Although organic limnic peat extends almost to the bottom, little or no tree pollen is present below 5.2 meters, so as to suggest that forests did not invade the region for some time after the initiation of the sediments. Birch is poorly represented at higher elevations in the Matanuska Valley and on the tableland between the Talkeetna and Chugach mountains. The five sections between mileposts 88 and 158 were taken from swampy lake margins at elevations of 2500 feet. Spruce is recorded to practically 100 per cent in all of these columns, while occasional pollen grains of pine and mountain hemlock undoubtedly were carried inland from the coast. A 3.9-meter section at milepost 158 lacks tree pollen in the lower 0.9 meter, suggesting the absence of forests within range of pollen dispersal for some time after deposition had begun.

### POSTGLACIAL CHRONOLOGY AND CLIMATE

The term "postglacial" is used rather loosely here in that it refers to the time since each sedimentary site was freed of ice and the pollen-bearing sediments began to accumulate. Obviously the time that sedimentation began must have varied from site to site, or at least in the several regions represented. Also the time when vegetation first appeared within range of pollen dispersal to each respective site may have varied considerably. Inasmuch as the glacial chronology of Alaska and of the Yukon Territory is poorly defined and there is evidence that neither all of Alaska nor the Yukon was glaciated during Wisconsin time, the usage of "postglacial" does not imply post-Mankato or post-Late Wisconsin as it usually does farther south in Canada and the United States. It seems probable, however, that the sections described in this study are composed of sediments that have accumulated since the glacial retreat late in the Wisconsin age.

In general the depth of the muskegs developed under nonaquatic conditions decreases with an increase in latitude. This reflects the less favorable growing conditions for the principal peat-forming plants, caused by the shorter growing season. Terraces of stratified silt and sand 200 feet above the present level of the Tanana River indicate the height of the river during deglaciation, and peat sections obtained from sites at this or lower elevations are necessarily postglacial (Brooks,

1900). The vast unglaciated area of interior Alaska provides the possibility of continuous peat deposition since a time before retreat of the late Wisconsin glaciers in the Alaska Range to the south and the Brooks Range to the north. Inasmuch as muskeg grades into tundra northward and both types of vegetative cover result in organic deposition, it is conceivable that on sites beyond and above the direct and indirect physiographic effects of glaciation on this interior plateau, peat deposition may have been initiated and continued regardless of glacial conditions in the mountain ranges. There is no reason to assume that the climate during the Wisconsin glaciation was necessarily more severe than it is today on the Arctic plain north of the Brooks Range which supports luxuriant tundra vegetation. The persistence of plants in unglaciated refugia in the Yukon and Alaska as pointed out by Hultén (1937) supports this view. In fact, it has been suggested that the climate of Alaska was warmer than normal during the ice age and at no time colder than the present (Pauly, 1952). The abundance of vegetal remains, including trees, and strata of interglacial peat in the mucks and silts in the Fairbanks region, as well as in other parts of Alaska (Taber, 1943; Tuck, 1940), would indicate that with each glacial age the vegetation of interior Alaska and other ice-free areas in the Yukon, British Columbia, and Alberta was not entirely obliterated. It is hardly conceivable that during each interglacial age vegetation migrated from beyond the southern limits of glaciation on the continent all the way back to interior Alaska and west-central Yukon. The abundance of lodgepole pine pollen in the lowest levels of peat sections in northeastern British Columbia and western Alberta causes the author to believe that late during the Wisconsin glaciation there were ice-free areas that supported forests of lodgepole and spruce (Hansen, 1949*a*, 1949*b*, 1950*a*, 1952). The position of the "Altamont" (Mankato) moraine, 70 miles east of Edmonton, further supports this evidence (Bretz, 1943).

The author does not presume, however, that any of the peat sections of this study antedate the beginning of the final recession of the Late Wisconsin glacier in this region. As stated above, the sections from the Tanana River valley are too close to the river to believe that the terrain upon which they lie was undisturbed or not indirectly modified by late Wisconsin glaciation in the Alaska Range not far to the south (Capps, 1931). The other sections were taken from sites that were either

overrun by ice and/or modified by meltwater. Many of the peat deposits have accumulated in standing water that was ponded by glaciofluvial processes resulting directly from deglaciation. The author does believe, upon the basis of the evidence, that the ages of the pollen-bearing sections do differ considerably. It is interesting and significant that in sections along the Alaska Highway in the Yukon and Alaska the shallower muskegs contain forest tree pollen in the lowest horizons and even in the upper part of the inorganic silt, while in the deeper sections that have been formed in ponds and lakes, tree pollen is largely or entirely absent in the lower portions of the columns, although aquatic plant pollen may occur almost to the bottom. The two thickest sections in the Yukon, 4.9 and 5.5 meters at mileposts 859 and 872 respectively, have more than a meter of the lowest portions devoid of tree pollen. Likewise, two sections along the Slana-Tok Highway, 3.0 and 2.0 deep, reveal only 1.6 and 0.9 meters respectively of tree pollen-bearing sediments. The same is true along the Palmer-Glenn Highway east of Anchorage, where the thickest section of this study, 6.9 meters, contains arboreal pollen down only to 5.2 meters, while a column 4.5 meters thick contains tree pollen only to the 2.6 meter horizon. The presence of nonarboreal pollen denotes that the accumulating sediments were receptive to and favorable for the preservation of pollen, while the absence of tree pollen suggests that forests were absent within range of pollen dispersal for some time after sedimentation began. Apparently conditions were favorable for the existence of plants in the region. Pollen from forests that may have existed in the unglaciated interior may have been prevented from reaching these sites because of prevailing westerly winds.

In the typical muskeg the presence of ericad and spruce pollen down to the underlying inorganic material suggests that forests and other vegetation were in existence when organic deposition in these muskegs began. In a few muskegs on flat terrain without early aquatic stages, spruce pollen does not occur all the way to the bottom. This also denotes the absence of forests in adjacent areas unless winds prevented pollen from reaching the site of the sediments. The age then of the lowest pollen-bearing sediments is uncertain.

Rate of peat deposition in this region must be exceedingly slow because of the rigorous climate and the short growing season. The thickness of the typical muskeg deposit increases

southeastward along the Alaska Highway and southward still farther to the vicinity of Edmonton, Alberta, which is the approximate limit in Alberta of what the author considers to be true muskeg (Hansen, 1949*a*, 1949*b*, 1950*a*, 1952). In the latter region the muskeg depth above the residual blue clay formed by gleization of waterlogged glacial drift averages about 3 meters as compared to the average of 0.5 meter depth of muskegs between the Yukon border and Fairbanks (table 1). The thickest section obtained from a muskeg along the Alaska Highway is 7.0 meters at milepost 253 south of Fort Nelson (Hansen, 1950*a*). It would seem that the rate of deposition under the conditions at this latitude is much slower than farther south where the climate is more favorable, although the relative ages of the muskegs in the two regions must be considered.

Several yardsticks may be used in estimating the age of the peat sections of this study based upon data derived directly from peat in this region. Capps (1931) studied the relationships between the development of adventitious roots of spruce and progressive rise of the permafrost table in a living *Sphagnum* moss deposit resting on fresh till about eight miles beyond the terminum of Russell Glacier. This site is located in an exposure cut by the White River near the Yukon-Alaska border. Calculations based upon annual ring counts indicate that the peat has accumulated at a rate of one foot in 200 years, which when applied to the 39-foot thick deposit denotes an age of about 7800 years. This rate applied to the muskegs in the Tanana River valley would indicate an age of only a few hundred years. It is possible that the White River deposit represents optimum conditions for muskeg development, as it can be readily demonstrated that the rate of peat accumulation may vary as much as 9 to 1 in a single region (Hansen, 1947). This rate applied to the thickest section of this study, 15 miles east of Anchorage, would imply an age of about 4600 years. While the White River peat site undoubtedly was covered by late Wisconsin ice, the cutting of a gorge 70 feet or more deep may have confined its meltwater to the gorge, thereby preventing disturbance of the peat site early in postglacial time, while in the Tanana River valley disturbance of the present muskeg sites by meltwater for a much longer time may be reflected in the shallower muskegs. A similar interpretation may be applied to the deeper sections of this study that have developed in valleys carrying meltwater from Wisconsin glaciers.

A second chronological yardstick available is the radiocarbon date of peat from the Anchorage area. A sample of peat from the lower foot of an 8-foot exposure at the north end of the Anchorage International Airport has been dated at about 5000 years (Kulp, Feely, and Tryon, 1951). This figure implies an accumulation rate of more than 600 years per foot, which when applied to the thickest section of this study near Anchorage denotes an age of more than 13,000 years. Another radiocarbon date of significance is that of about 3800 years for peat at a depth of five feet, near Boulder Point, Kenai Peninsula (Kulp, et al., 1952). Peat accumulation here probably has occurred under climatic conditions somewhat similar to those in the Anchorage area. The rate of about 800 years per foot would indicate almost 18,000 years for the deep section near Anchorage. Climatic conditions for peat accumulation would seem to be more favorable today in the Anchorage region than in the interior. Further study of the White River peat exposure, including radiocarbon techniques and pollen analysis, would be extremely significant in that it would provide a valuable index for the chronology of postglacial time in Alaska and the Yukon. Another radiocarbon date of significance is that of 14,300 ± 600 years determined from lignitic peat exposed along the Eagle River north of Anchorage, which is believed to be overlain by Late Wisconsin till, and appears to have been deposited during the Cary-Mankato interglacial. This figure and Capps' computations suggest that the sites represented were deglaciated as early as or earlier than the northern Lake States as denoted by radiocarbon dates of peat and wood from that region (Flint and Deevey, 1951). It becomes very evident that a series of carbon-14 dates for Alaska peats is essential for better approximation of their ages and application to the forest record in this region. Heusser (1952), in a pollen and stratigraphic study of 17 peat sections in southeastern Alaska from Ketchikan to Juneau, sets a date of about 8000 years for the older and deeper muskegs. These muskegs vary in depth from a few decimeters to 6 meters.

There seems to be little or no evidence for climatic trends in the pollen profiles, except perhaps a general amelioration throughout the time represented. This is expressed by a rapid expansion of lodgepole to its maximum in many of the Yukon sections, although some of them reveal a preponderance of pine throughout. The thickness of the peat through which lodgepole

is predominantly recorded after attaining its maximum varies considerably. At milepost 647 it is recorded to 80 per cent or more in the upper 3 meters of a 5.6-meter section, while in other thick sections it is preponderantly recorded in only the upper few decimeters. This difference can be assumed to be caused by local fluctuations in the relative abundance of spruce and pine as well as a variation in the rate of peat deposition. The occurrence of two volcanic glass layers within the period of pine predominance, however, indicates general synchronism in the several profiles. In many sections a slight increase in pine in the upper decimeter or two may reflect the influence of fire since the advent of white man. It is not believed that the warmer and drier climate of the past 200 years as indicated by glacial retreat in southeastern Alaska and dated by dendrochronological studies is reflected by this pine increase (Lawrence, 1950). The upper decimeter of pine increase probably constitutes less than 200 years, while the total period of pine maximum is much greater than 200 years. The sections in Alaska do not manifest climatic trends and apparently the postglacial climate has not been sufficiently warm to favor migration of pine into the interior. While the author believes that most of the deeper pollen-bearing sediments extend back to the postglacial thermal maximum so well shown elsewhere, there seems to be no evidence that the influence of this warm, dry interval extended this far north. The abundance of birch in the Anchorage region may reflect both fire and the absence of perennially frozen ground (Taber, 1943).

The pollen assemblages of this study indicate that the present range of lodgepole pine does not differ significantly from that during any of the time represented by the pollen profiles. The presence of appreciable pine pollen in the lowest levels of the Yukon sections denotes that pine was prevalent in the region when the earliest pollen-bearing sediments were deposited, and that it probably persisted during the late Wisconsin glaciation in refugia not far removed from the sites of those sediments. It may have migrated northwestward from ice-free areas in western Alberta and/or northeastern British Columbia or from refugia in west-central Yukon. The former hypothesis is supported by the lodgepole pine pollen record in muskegs in Alberta and British Columbia (Hansen, 1950*a*, 1952). The absence of

pine wood or other macroscopic remains of this species in the Pleistocene mucks and silts in the Fairbanks region suggests that it did not migrate into interior Alaska during Pleistocene time (Chaney and Mason, 1936). The occurrence of a few pine pollen grains observed in a preliminary study of these sediments by the author may mean merely the same as their occurrence in the postglacial sediments of this study, namely, that pine pollen was carried into the area from distant sites. The early postglacial appearance of lodgepole pine in coastal southeastern Alaska as expressed by Heusser's studies may have resulted from its migration down the Stikine River valley from interior British Columbia, or over White Pass from the Bennett Lake area into the Lynn Canal drainage basin. A northward coastal migration of lodgepole from centers of its persistence south of the glacial border in Washington seems less probable.

### References


Bostock, H. S., 1948. Physiography of the Canadian Cordillera, with special reference to the area north of the fifty-fifth parallel: Canada Geol. Survey Mem. 247.

Bretz, J Harlen, 1943. Keewatin end moraines in Alberta, Canada: Geol. Soc. America Bull., vol. 54, pp. 31-52.

Brooks, A. H., 1900. A reconnaissance in the White and Tanana River basins, Alaska, in 1898: U. S. Geol. Survey 20th Ann. Rept., part 7.

Capps, S. R., 1916. The Chisana-White River district, Alaska: U. S. Geol. Survey Bull. 630.

———, 1931. Glaciation in Alaska: U. S. Geol. Survey Prof. Paper 170.

Chaney, R. W., and Mason, H. D., 1936. A Pleistocene flora from Fairbanks, Alaska: Am. Mus. Novitates 887.

Denny, C. S., 1952. Late Quaternary geology and frost phenomena along Alaska Highway, northern British Columbia and southeastern Yukon: Geol. Soc. America Bull., vol. 63, pp. 883-922.

Flint, R. F., 1947. Glacial geology and the Pleistocene epoch, John Wiley & Sons, Inc., New York.

———, et al., 1945. Glacial map of North America: Geol. Soc. America Special Paper 60.

———, and Deevey, E. S., Jr., 1951. Radiocarbon dating of late-Pleistocene events: Am. Jour. Sci., vol. 249, pp. 257-300.

Halliday, W. E. D., 1937. A forest classification of Canada: Canada Dept. Mines and Resources, Forest Survey Bull. 89.

———, and Brown, A. W. A., 1943. The distribution of some important forest trees in Canada: Ecology, vol. 24, pp. 353-374.

Hansen, H. P., 1947. Postglacial forest succession, climate, and chronology in the Pacific Northwest: Am. Philos. Soc. Trans., new ser., vol. 37, part 1.

———, 1949*a*. Postglacial forests in south central Alberta, Canada: Am. Jour. Botany, vol. 36, pp. 54-65.

———, 1949*b*. Postglacial forests in west central Alberta, Canada: Torrey Bot. Club Bull., vol. 76, pp. 278-289.

———, 1950*a*. Postglacial forests along the Alaska Highway in British Columbia: Am. Philos. Soc. Proc., vol. 94, pp. 411-421.

———, 1950*b*. Pollen analysis of three bogs on Vancouver Island, Canada: Jour. Ecology, vol. 38, pp. 270-276.

———, 1952. Postglacial forests in the Grande Prairie-Lesser Slave Lake region of Alberta, Canada: Ecology, vol. 33, pp. 31-41.

Harlow, W. M., and Harrar, E. S., 1941. Textbook of dendrology, McGraw-Hill Book Company, New York.

Heintzelman, F. B., 1951. Forestry in Alaska: Alaskan Science Conf. Bull. Proc., Nat. Res. Council 122, pp. 126-133.

Heusser, C. J., 1952. Pollen profiles from southeastern Alaska: Ecology Mon. 22, pp. 331-352.

Hultén, Eric, 1937. Outline of the history of arctic and boreal biota during the Quaternary period, Bokförlags aktie bolaget Thule, Stockholm.

Kerr, F. A., 1936. Physiography of the cordilleran region of northern British Columbia and adjacent areas: Royal Soc. Canada Trans., 3d ser., vol. 30, sec. 4, pp. 137-154.

Kulp, J. L., Feely, H. W., and Tryon, L. E., 1951. Lamont natural radiocarbon measurements, 1: Science, vol. 114, pp. 565-568.

———, Tryon, L. E., Eckelman, W. R., and Snell, W. A., 1952. Lamont natural radiocarbon measurements, 2: Science, vol. 116, pp. 409-414.

Lawrence, D. B., 1950. Glacier fluctuation for six centuries in southeastern Alaska and its relation to solar activity: Geog. Rev., vol. 40, pp. 191-223.

Lutz, H. J., 1951. Ecological effects of forest fires in the interior of Alaska: Alaskan Science Conf. Proc. Bull., Nat. Res. Council 122, p. 21.

Moss, E. H., 1949. Natural pine hybrids in Alberta: Canadian Jour. Research, vol. 27, pp. 218-229.

Munns, E. N., 1938. The distribution of important forest trees of the United States: U. S. Dept. Agr. Misc. Bull. 287.

Pauly, K. A., 1952. The cause of the great Ice Age: Sci. Monthly, vol. 75, pp. 89-98.

Raup, H. M., 1935. Botanical investigations in Wood Buffalo Park: Canada Nat. Mus. Bull. 74.

———, 1946. Phytogeographic studies in the Athabaska-Great Slave Lake region: II: Jour. Arnold Arboretum. vol. 27, pp. 1-85.

Stoeckler, E. G., 1949. Identification and evaluation of Alaskan vegetation from airphotos with reference to soil, moisture and permafrost conditions: Army Engineers Corps, Field Operation Branch, Permafrost Div., St. Paul, Minn.

Taber, Stephen, 1943. Perennially frozen ground in Alaska; its origin and history: Geol. Soc. America Bull., vol. 54, pp. 1433-1548.

Taylor, R. F., 1950. Pocket guide to Alaska trees, U. S. Dept. Agr. Forestry Service Handbook 5, Washington.

Tuck, Ralph, 1940. Origin of the muck deposits at Fairbanks, Alaska: Geol. Soc. America Bull., vol. 51, pp. 1295-1310.

Wenner, Carl-Gösta, 1947. Pollen diagrams from Labrador: Geog. Annaler, vol. 29, pp. 137-174.

Oregon State College
Corvallis, Oregon

[American Journal of Science, Vol. 251, July 1953, Pp. 543-546]

# AGE OF INTERTIDAL TREE STUMPS AT ROBINHOOD, MAINE

W. H. BRADLEY *

ABSTRACT. Well-preserved stumps of eastern white pine whose age has been determined by carbon 14 analysis to be about 4000 years old, were found between tides at Robinhood, Maine. The stumps are 3 to 5 feet below mean high tide and indicate that when the trees lived mean high tide was probably 8 or 9 feet below its present level. These stumps and the peat beds at the Boston fish weir site, which are of comparable age, suggest a marked eustatic lowering of sea level immediately after the Climatic Optimum.

CHANGES of sea level are significant events in geologic history, and all means possible should be used to date them. In the summers of 1951 and 1952, while studying the microstratigraphy and sedimentation regimen of a large tidal flat on the Maine coast, I had an opportunity to observe, between high and low tide lines, the rooted stumps of trees in place. At one locality, about 13 miles southeast of Bath, I mapped their positions and collected specimens of wood from the largest stump. This wood has been identified by R. W. Brown of the U. S. Geological Survey as eastern white pine (*Pinus strobus*). Its age, determined by L. J. Kulp of Columbia University by carbon 14 analysis, is 4150 years ± 200.

The locality studied is on Georgetown Island near the village of Robinhood, situated on a long crooked arm of the sea that connects the Kennebec River with the Sheepscot River. North of Robinhood, across Riggs Cove, are two nearly circular hills, the Knubble and the Little Knubble. Near the center of a short, gravelly tombolo connecting these two hills, ancient trees grew on a site now 3.5 to 5 feet below mean high tide. The tombolo is of glacial origin and hence long antedates the trees (plate 1).

The mean tidal range at this site is a little less than 10 feet and, as a result of constriction of the channel between the Knubble and the Little Knubble, rather strong currents move across the tombolo, preventing sand and mud from accumulating on the gravel and stumps. Miss Sayde Campbell, owner of the Knubbles, reported that these stumps were known to her

* Publication authorized by the Director, U. S. Geological Survey.

grandmother and that probably they have been exposed, much as they are today, for at least 75 years.

That the trees grew here is clearly evident from the fact that many of the roots can be traced into the gravel. Some roots lace in and out of the gravel, not only between the stumps but for several yards beyond them. This is especially true west and north of the stumps. The roots are embedded in a black peat or soil that has worked its way down among the cobbles and fills hollows in the tombolo surface. Locally this black peaty material, which contains many small rectangular fragments of wood, is as much as a foot thick below the base of the lowest stump. All roots embedded in this material are soft and spongy, as though deeply decayed. Roots and stumps that rise above the peaty material, however, are quite sound and appear as though they could not have been there many years. Although thoroughly water-soaked, the wood chopped like perfectly sound wood and was obviously uniform to the bottom of the kerf, from which the samples were taken. The wood was bright turmeric yellow while wet and had a strong, disagreeable odor that could not be identified. After drying a few days in the sun, the wood lost both its yellow color and its bad odor.

After the wood chips were dried, all surface material of the chips was cut away so as to minimize the danger of intrusion by younger organisms, which would reduce the age as determined by carbon 14 analysis. None of the peaty material around the roots was collected for checking age determinations because of the high probability that both the grass (*Spartina alterniflora*) now growing close by and a variety of small marine animals had invaded the peaty material.

At a few other places along the intertidal zone around the two Knubbles, smaller stumps and parts of trees were found that may be the age equivalents of the stumps and roots described. Each of these other trees, however, had peculiarities that make the reliability of this inference doubtful. On the other hand, in 1951 and again in 1952, fairly well-preserved stumps were observed, apparently in positions where they had grown, on Prouts Neck, a few miles southwest of Portland. These stumps project through the sand on a steep, narrow beach that extends along the east side of the Scarborough River inlet, close to Western Beach. As at the Robinhood locality, the stumps are only a few feet below the mean high tide

PLATE 1

View eastward along the tombolo toward Little Knubble, showing stumps of trees in a tide pool. The approximate position of mean high tide is indicated by dashed line.

line in an area where tidal scour prevents the accumulation of sand in quantity.

Other drowned forests along the North Atlantic coast (Bay of Fundy, N. S.; Grand Pre, on an arm of the Bay of Fundy; Odiorne Point near Portsmouth, N. H.; and Provincetown, Cape Cod) have been known for many years. According to the description given by Lyon and Goldthwait (1934) the stumps at Odiorne Point and at Provincetown occupy about the same intertidal position as those at Robinhood, Maine. Possibly they are contemporaneous.

The tree stumps at the Robinhood locality obviously grew when sea level was at least 7 feet lower than at present, for the base of the lowest stump is about 5 feet below the mean high tide line and rests on about a foot of peaty soil that contains many wood fragments. Much more likely, however, sea level was some 8 or 9 feet below its present mean. This is inferred from the habit of the white pine and other conifers that grow in dense stands along the coast today. Rarely are the bases of these living trees less than 2 or 3 feet above the mean high tide.

In the speculation about former sea levels I have assumed that the changes were all eustatic and that they were caused by climatic changes, which either released water from the polar ice caps, which resulted in higher sea levels, or locked up water in expanding polar ice caps, which resulted in lowered sea levels. Thus presumably during the Climatic Optimum sea level was several feet higher than it is today. This assumption may be an unwarranted simplification, but it provides a simple, and not unreasonable, framework within which to state the significance of the Robinhood tree stumps.

The ancient tree stumps at Robinhood indicate that about 4000 years ago mean sea level was at least 7 to 9 feet lower than it is today. The Boston fish weir, whose age is somewhere between 3851 $\pm$390, and 5717 $\pm$ 500 years old (Arnold and Libby, 1951, p. 113) suggests that sea level may have been as much as 29 feet below its present position (Johnson, 1942, p. 161). If the Climatic Optimum occurred sometime between 4000 and 8000 years ago, the Robinhood tree stumps and the Boston fish weir suggest that there may have been a significant glacial readvance at the end of the Climatic Optimum to account for a sizeable drop of sea level.

Presumably, the ancient trees at Robinhood were killed by the rise of sea level after this inferred cold interval because a discontinuous layer of peat (presumably all *Spartina*) 9 to 12 inches thick apparently once covered, or surrounded, the tree stumps. This peat has been partly eroded by tidal scour or ice plucking, and its eroded edges are nearly vertical. *Spartina* now grows on most of its surface (plate 1). I infer that this irregular layer of peat grew when sea level was still low enough to prevent normal tides from overtopping the tombolo. Clearly this peat and its present growth of grass are today in a precarious position—one subject more to erosion than to growth and extension.

Lyon and Goldthwait (1934, p. 611) suggest that the drowned stumps at Odiorne Point were "first buried by salt marsh and exposed in comparatively recent time."

My postulate that the Robinhood trees indicate a significant drop in sea level at the end of the Climatic Optimum is obviously based on slender evidence—one point on the curve, one might say. Nevertheless, the material for the carbon 14 age determination appears to be good and the evidence at the site rather simple and clear. I have not tried to relate my observations and speculation to possible regional movement of the coast following the ice age because I frankly do not know how to get hold of the problem.

Perhaps these observations and my speculation will serve to encourage others to get more and better "points on the curve" to the end that we may someday gain a full understanding of that fascinating part of geologic history which must be so closely linked with man's early history.

## References

Lyon, C. J., and Goldthwait, J. W., 1934. An attempt to cross-date trees in drowned forests: Geog. Rev., vol. 24, pp. 610-611.

Arnold, J. R., and Libby, W. F., 1951. Radiocarbon dates: Science, vol. 113, p. 113.

Johnson, Fredrick, et al., 1942. The Boylston Street fish weir: Robert S. Peabody Foundation for Archeology Papers, vol. 2.

U. S. Geological Survey
Washington, D. C.

[American Journal of Science, Vol. 251, July 1953, Pp. 547-548]

# DISCUSSION

## FOSSIL COMMUNITIES AND ASSEMBLAGES

With the merging of paleontology and ecology in the subject of paleoecology a confused terminology has arisen, partly ecological, partly paleontological. It is to be hoped that the proposed *Treatise on Marine Ecology and Paleoecology* will rectify this confusion and establish a terminology acceptable to the majority of ecologists and paleontologists; a terminology, moreover, which will involve the use of a minimum of technical jargon. Boucot in this Journal (Vol. 251, January 1953, p. 26) has suggested that "life assemblage" and "death assemblage" should be used as the paleontological equivalents of "biocoenosis" and "thanatocoenosis" respectively. While Boucot makes a welcome departure from the prevailing tendency to establish a jargon by latinizing or hellenizing short, understandable, English phrases, it is unfortunate that his terms are ambiguous. To avoid this ambiguity one would have to write of a "fossil life assemblage" and a "fossil death assemblage," the former seemingly paradoxical, the latter repetitive.

Ecologists use "community" in the sense of an ecologically related association of organisms, especially marine bottom animals (see Jones, 1950). The term is of obvious application to paleoecology. A "fossil community" can be considered as an association of fossils ecologically related among themselves and to their containing deposit, while a "fossil assemblage" can be used, without ecological implications, of an association of fossils, as is normal practice. A "fossil assemblage" may consist of one or several fossil communities; it may be wholly composed of fossils ecologically foreign to the containing deposit; or it may consist of a mixture of foreign and indigenous fossils.

The term "thanatocoenosis" because of its varied usage is avoided (cf. Wasmund, 1926; Twenhofel, 1939, p. 154; Wells, 1947, p. 119; Ellison, 1951, p. 214). It might perhaps be added that the study of fossil assemblages has been termed "paleothanatology" (Morishima, 1948, p. 111).

### References

Boucot, A. J., 1953, Life and death assemblages among fossils: Am. Jour. Sci., vol. 251, p. 25-40.

Ellison, S. P., 1951, Microfossils as environment indicators in marine shales: Jour. Sedimentary Petrology, vol. 251, p. 214-225.

Jones, N. S., 1950, Marine bottom communities: Cambridge Biol. Rev., vol. 25, p. 283-313.

Morishima, M., 1948, Foraminiferal thanatocoenoses of Ago Bay, Kii Peninsula, Japan: Comm. on Treatise on Marine Ecology and Paleoecology Rept. 8, Nat. Research Council, Washington.

Twenhofel, W. H., 1949, Principles of sedimentation, McGraw-Hill Book Company, New York.

Wasmund, E., 1926, Biocoenose und Thanatocoenose: Archiv Hydrobiologie, vol. 17, p. 1-116.

Wells, J. W., 1947, Provisional paleoecological analysis of the Devonian rocks of the Columbus region: Ohio Jour. Sci., vol. 47, p. 119-126.

GRANT INSTITUTE OF GEOLOGY
UNIVERSITY OF EDINBURGH
SCOTLAND

GORDON CRAIG

## REVIEWS

*Probleme der Naturwissenschaften* (*Erläutert am Begriff der Mineralart*); by PAUL NIGGLI. Pp. xi, 240; 99 figs. Basel, 1949 (Verlag Birkhäuser, 18.50 Swiss francs).—This book presents a very interesting and authoritative review of modern methods of mineralogical investigation as well as the results of that science. Its meat is sandwiched between lengthy preliminary remarks of a philosophical nature and a brief general conclusion which again carries a philosophical flavor. The main contents of the book form a pleasing and judicious selection of materials which a philosopher ought to know; an account is given of crystal structure and crystal types, of the technical meaning of symmetry, of genotypes and phenotypes, and of the effects of substitutions on crystal structures. These parts are eminently worth serious attention, even by the general reader who may find some of the discussion difficult.

The peripheral portions of the book cannot escape altogether the criticism of being unsystematic, even sometimes unprincipled tracts of philosophic writing. Here one finds rather indiscriminate references to many philosophers of different points of view without evidence of a unifying effort on the part of the author. The language is frequently difficult, as for instance, in the characterization of the two chief methods of scientific inquiry, which are "(α) die *abstrakt-generalisierende* oder *atektonisch-imperative* bzw. *ursächlich-erklärende*, die andere (β) die *vergleichend-systematische* oder *tektonisch-normative* bzw. *urbildlich-erläuternde*." Despite such faults the book has philosphical importance inasmuch as it contains a biographical statement of the points of view which a successful scientist has found significant in his own investigations.

HENRY MARGENAU

*Water: A Study of Its Properties, Its Constitution, Its Circulation on the Earth, and Its Utilization by Man*; by CYRIL S. FOX. Pp. xxvii, 148; 4 figs., 25 plates. New York, 1952 (Philosophical Library, $8.75).—This little book is the introductory volume of a series of books which the Technical Press, Ltd., has in preparation, or has already published, "to meet the current need for information on the various aspects of water supply." The titles of some of the other volumes of the series are *The Geology of Water Supply, Water Treatment,* and *Purification and Disposal of Sewage.* Some of these volumes have already been published—for example, Sir Cyril's *Geology of Water Supply.* The author aims at filling a need that has long existed, namely, to treat of water as a subject for coordinated study, rather than as a part of some other discipline such as hydraulics, meteorology, engineering, or geology.

Sir Cyril divides his book into a Preface, Foreword, and Introduction—27 pages; Part 1, "The Natural History of Water," with chapters on the constitution, distribution, and circulation of water—68 pages; Part 2, "The Work Done by Water," with chapters on erosion, action of underground water, and deposition of sediments—53 pages; and Part 3, "The Utilization of Water," with chapters on general considerations, water supply, engineering, and concluding remarks—21 pages.

The photographs, 25 in number, are excellent and very well reproduced, but there are only 4 other illustrations. Much of the factual material would be found in any good textbook on physical geology, or in U. S. Geological Survey Bulletin 770, "Data of Geochemistry," and it cannot be said that the book adds much to our knowledge of water. No new material is presented except the statements on the radioactive water in Bikini.

Some of the statements made will be rather startling to geologists and engineers engaged in water-resources development or appraisal. For example, the assertion on page 128 that "Many irrigation engineers can estimate very closely the flow of a moderate-sized stream by a careful scrutiny" would, if true, eliminate the need for a rather large and active program of stream gaging now being carried on by federal and state governments and by private enterprise. Most hydraulic engineers would deny the ability to estimate stream discharge within 25 to 50 per cent. The further statement that "for these larger schemes accurate determinations are necessary, perhaps for two or three years to secure low and flood water discharges" suggests that the author's experience is limited to areas where conditions of precipitation are much more uniform than any encountered in the United States, where a minimum of 10 to 20 years of record are considered necessary for sound planning and design of water facilities. Another statement, "Among civil engineers

it is a common rule that rainfall is disposed of in equal proportions as runoff flow, loss of evaporation and loss by percolation," is at variance with the findings of W. G. Hoyt and others, as reported in Water-Supply Paper 772, "Rainfall and Runoff."

One particularly serious fault in the book is the author's tendency to draw sweeping generalizations on the basis of averages which he later demonstrates to be false. On page 37 Sir Cyril states, "It is computed that there is a temperature increase of 1°C. for every 100 ft. descent in depth. If this is uniform the temperature at a depth of 2,500 feet . . . is 25°C. plus the surface temperature. . . . " But on page 96 he demonstrates that the data are meager and such data as are available rarely agree with the assumed average.

The objective of this book is one with which the reviewer is in complete sympathy, and it is hoped that in spite of its shortcomings this particular volume will serve to stimulate others to "draw attention to the immense importance of the control and use of water by first understanding the practical aspects of the material itself."

A. NELSON SAYRE

*Numerical Mathematical Analysis*, 2d ed.; by JAMES B. SCARBOROUGH. Pp. xviii, 511; 51 figs. Baltimore, 1950 (The Johns Hopkins Press, $6.00).—The first edition of this book appeared in 1930. Since then the development of powerful rapid calculators has made possible numerical attacks upon problems that were formerly considered too laborious. This has stimulated a revival of interest in the processes of numerical mathematical analysis. The new edition reflects this recent development. As might be expected, the most significant changes are in chapters dealing with differential equations. The chapter on the numerical solution of ordinary differential equations is greatly expanded, and entirely new chapters on the numerical solution of partial differential equations and of integral equations have been added. Thus the usefulness of the book has been increased.

The clarity of the presentation is on a high level. It stresses the principles and processes involved, and evaluates the limitations of practical methods both by analytical treatment and numerical examples. The valuable comments that accompany many of these examples give ample evidence of the author's insight and experience.

DIRK BROUWER

*The Heavens Above. A Rationale of Astronomy*; by J. B. SIDGWICK. American edition prepared by WARREN K. GREEN. Pp. xv, 333; 56 figs., 20 plates, New York, 1950 (Oxford University Press, $4.00).—The more spectacular results of modern astronomi-

cal research have been accessible to the reading public through numerous popular books, some of which were among the bestsellers. The authors of these books have not always made it their duty to make clear the basic principles, the trains of reasoning, and the observations upon which this edifice was built. The purpose of this book is to furnish a systematic demonstration of the more complex facts of astronomy.

In the first part of the book, Quantity—a geometric picture, the spatial arrangement of the astronomical bodies in the universe is presented. The logical process that leads to the determination of astronomical distances from the nearest neighbors of the earth to the most distant galaxies is presented with attention to the margin of uncertainty that affects the successive extrapolations. It is necessary occasionally to make use of principles and facts presented in the second part, Quality—the nature of things. This is done with little damage to the continuity of the presentation.

The purpose that prompted the author to write this book is precisely the same as that of many elementary courses on the college level. For this reason the book is eminently suitable to serve as a textbook for such courses. Advantages over some standard texts are its refreshing style and its pleasing arrangement.

Among the few less satisfactory parts of the book is the introduction to spectroscopy. Too many details about energy levels and series of spectral lines are presented in too small a space on insufficient foundation. A simplification of this part would be an improvement. When dealing with planetary atmospheres, the notion that the giant planets have tremendously deep atmospheres is presented as an established fact, while conclusive evidence to the contrary is now available. Over-cautiousness led the author to question whether the galactic star system is coextensive with the system of open clusters or with the system of globular clusters. This question has been definitely resolved in favor of the latter alternative. These imperfections do not seriously affect the otherwise excellent quality of the presentation. DIRK BROUWER

## PUBLICATIONS RECENTLY RECEIVED

Organic Syntheses, Vol. 32; R. T. Arnold, editor-in-chief. New York and London, 1952 (John Wiley & Sons, Inc., and Chapman & Hall, Ltd., $3.50).

Soil Microbiology; by S. A. Waksman. New York and London, 1952 (John Wiley & Sons, Inc., and Chapman & Hall, Ltd., $6.00).

A University Textbook of Physics, Vol. III, Heat; by J. H. Awbery. London, 1952 (Charles Griffin & Co., Ltd., $5.50 in U.S.A.; 36s., plus 1s. 6d overseas postage).

Amenability of Certain Kansas Clays to Alumina Extraction by the Lime-sinter Process; by E. D. Kinney. Kansas State Geological Survey Bulletin 96, Part 7. Lawrence, 1952 (University of Kansas).

U. S. Geological Survey Water-supply Papers as follows: 1110-D, Sedimentation Rates in Small Reservoirs in the Little Colorado River Basin; by C. F. Hains, D. M. Van Sickle, and H. V. Peterson ($ .40). 1134-B, Floods of July 18, 1942 in North-central Pennsylvania; by W. S. Eisenlohr, Jr. ($ .75). 1137-B, Floods of 1950 in the Red River of the North and Winnipeg River Basins ($1.00). 1137-C, Flood of August 1950 in the Waimea Area, Kauai, Hawaii; by R. K. Chun ($ .15). 1139, Kansas-Missouri Floods of July, 1951 ($1.25). Surface Water Supply of the United States, 1949: 1141, Part 1, North Atlantic Slope Basins ($2.00); 1152, Part 12, Pacific Slope Basins in Washington and Upper Columbia River Basin ($1.00); 1153, Part 13, Snake River Basin ($ .60). Water Levels and Artesian Pressure in Observation Wells in the United States in 1949: 1156, Part 1, Northeastern States ($1.00); 1157, Part 2, Southeastern States ($ .55); 1160, Part 5, Northwestern States ($ .40). 1168, Water Levels and Artesian Pressure in Observation Wells in the United States in 1950, Part 4, South-Central States ($ .40). Surface Water Supply of the United States, 1950: 1175, Part 5, Hudson Bay and Upper Mississippi River Basin ($1.50); 1177, Part 7, Lower Mississippi River Basin ($1.00); 1178, Part 8, Western Gulf of Mexico Basins ($1.00); 1179, Part 9, Colorado River Basin ($1.00); 1180, Part 10, The Great Basin ($ .55); 1181, Part 11, Pacific Slope Basins in California ($1.00). Washington, 1952 (U. S. Government Printing Office).

U. S. Geological Survey Bulletins as follows: 972, Geology of the Canyon Ferry Quadrangle, Montana; by J. B. Mertie, Jr., R. P. Fischer, and S. W. Hobbs ($ .75). 973-B, The Coalwood Coal Field, Powder River County, Montana; by R. P. Bryson ($3.25, buckram). 974-E, The Eastern Front of the Bitterroot Range, Montana; by C. P. Ross ($1.00). 975-B, Geology of the Quicksilver Deposits of Canoas, Zacatecas, Mexico; by David Gallagher ($1.00). 979-B, Manganese Deposits of Southeastern Utah; by A. A. Baker, D. C. Duncan, and C. B. Hunt ($ .60). 982-A, Phosphatic Rocks in the Deer Creek-Wells Canyon Area, Idaho; by W. R. Lowell ($ .45). 983, Corundum Deposits of Montana; by S. E. Clabaugh. 985, Annotated Bibliography of North American Geology, 1950; by Marjorie Hooker ($1.00). 988-A, Geology of the Uravan Mineral Belt; by R. P. Fischer and L. S. Hilpert ($ .35). 988-B, Geologic Guides to Prospecting for Carnotite Deposits on Colorado Plateau; by D. B. Weir ($ .15). 991-A, Geophysical Abstracts 148, January-March 1952 (Nos. 13284-13547); by M. C. Rabbitt and S. T. Vesselowsky ($ .25). 991-B, Geophysical Abstracts 149, April-June 1952 (Nos. 13548-13802); by M. C. Rabbitt and S. T. Vesselowsky ($ .25). Washington, 1952 (U. S. Government Printing Office).

Essentials of Microwaves; by R. B. Muchmore. New York and London, 1952 (John Wiley & Sons, Inc., and Chapman & Hall, Ltd., $4.50).

An Introduction to Scientific Research; by E. B. Wilson, Jr. New York, 1952 (McGraw Hill Book Company, $6.00).

Mental Prodigies; by Fred Barlow. New York, 1952 (Philosophical Library, $4.75).

Development of the Guided Missile; by K. W. Gatland. New York, 1952 (Philosophical Library, $3.75).

VOL. 251 No. 7, 1953

# CONTENTS

PETROLOGICAL CALCULATIONS IN METASOMATIC PROCESSES
*Arie Poldervaart* 481

POSTGLACIAL FORESTS IN THE YUKON TERRITORY AND ALASKA
*Henry P. Hansen* 505

AGE OF INTERTIDAL TREE STUMPS AT ROBINHOOD, MAINE
*W. H. Bradley* 543

## DISCUSSION

Fossil Communities and Assemblages ..... *Gordon Craig* 547

## REVIEWS

Probleme der Naturwissenschaften (Erläutert am Begriff der Mineralart); by PAUL NIGGLI ..... *Henry Margenau* 548

Water: A Study of Its Properties, Its Constitution, Its Circulation on the Earth, and Its Utilization by Man; by CYRIL S. FOX ..*A. Nelson Sayre* 549

Numerical Mathematical Analysis, 2d ed.; by JAMES B. SCARBOROUGH
*Dirk Brouwer* 550

The Heavens Above. A Rationale of Astronomy; by J. B. SIDGWICK
*Dirk Brouwer* 550

PUBLICATIONS RECENTLY RECEIVED ..... 551

PAYNE & LANE, *Printers* — NEW HAVEN, CONN.